

Recitation 2G
February 10, 2005

1. Example 1.18, page 33 in the text. A test for a certain rare disease is assumed to be correct 95% of the time: if a person has the disease, the test results are positive with probability 0.95, and if the person does not have the disease, the test results are negative with probability 0.95. A random person drawn from a certain population has probability 0.001 of having the disease. Given that the person just tested positive, what is the probability of having the disease?
2. **Communication through a noisy channel.** Problem 1.27, page 59 in the text. A binary (0 or 1) symbol transmitted through a noisy communication channel is received incorrectly with probability e_0 and e_1 , respectively, as described in Figure 1. Errors in different symbol transmissions are independent.




Figure 1: Description of a binary channel

- (a) Suppose that the channel source transmits a 0 with probability p and transmits a 1 with probability $1 - p$. What is the probability that a randomly chosen symbol is received correctly?
 - (b) Suppose that the string of symbols 1011 is transmitted. What is the probability that all the symbols in the string are received correctly?
 - (c) In an effort to improve reliability, each symbol is transmitted three times and the received symbol is decoded by a majority rule. In other words, a 0 (or 1) is transmitted as a 000 (or 111, respectively), and it is decoded at the receiver as a 0 (or 1) if and only if the received three-symbol string contains at least two 0s (or 1s, respectively). What is the probability that a transmitted 0 is correctly decoded?
 - (d) Suppose that the source transmits a 0 with probability p and transmits a 1 with probability $1 - p$, and the scheme of part (c) is used. What is the probability that a 0 was transmitted given that the received string is 101?
3. A fair coin is tossed repeatedly. Show that the following two statements are equivalent:
 - (a) the outcomes of different tosses are independent,
 - (b) for any given finite sequence of heads and tails, the chance of this sequence occurring in the first m tosses is 2^{-m} , where m is the length of the sequence.

4. Show that the conditional independence of A and B given C neither implies, nor is implied by, the independence of A and B . For which events C is it the case that, for all A and B , the events A and B are independent if and only if they are conditionally independent given C ?