

Markov Processes – II

- **Readings:** Section 6.3

Lecture outline

- Markov process review
- Steady-State Behavior
- Birth-death processes

- Discrete state, discrete time, time-homogeneous
 - Transition probabilities p_{ij}
 - Markov property


- $r_{ij}(n) = P[X_n = j \mid X_0 = i]$

- Key recursion:

$$r_{ij}(n) = \sum_k r_{ik}(n-1)p_{kj}$$

Recurrent and transient states

- State i is **recurrent** if:
 - starting from i ,
 - and from wherever you can go,
 - there is a way of returning to i
- If not recurrent, called **transient**
- **Recurrent class:**
 - collection of recurrent states that “communicate” to each other and to no other state




$$P(X_1 = 2, X_2 = 6, X_3 = 7 \mid X_0 = 1) =$$

$$P(X_4 = 7 \mid X_0 = 2) =$$

Periodic states

- The states in a recurrent class are **periodic** if they can be grouped into $d > 1$ groups so that all transitions from one group lead to the next group



Steady-State Probabilities

- Do the $r_{ij}(n)$ converge to some π_j ? (independent of the initial state i)
- Yes, if:
 - recurrent states are all in a single class, and
 - no periodicity
- Start from key recursion

$$r_{ij}(n) = \sum_k r_{ik}(n-1)p_{kj}$$


- take the limit as $n \rightarrow \infty$

$$\pi_j = \sum_k \pi_k p_{kj}$$

- Additional equation:

$$\sum_j \pi_j = 1$$


Example





Visit frequency interpretation

$$\pi_j = \sum_k \pi_k p_{kj}$$

- (Long run) frequency of being in j : π_j
- Frequency of transitions $k \rightarrow j$: $\pi_k p_{kj}$
- Frequency of transitions into j : $\sum_k \pi_k p_{kj}$



Birth-death processes



$$\pi_i p_i = \pi_{i+1} q_{i+1}$$

- Special case: $p_i = p$ and $q_i = q$ for all i
 $\rho = p/q = \text{load factor}$

$$\pi_{i+1} = \pi_i \frac{p}{q} = \pi_i \rho$$

$$\pi_i = \pi_0 \rho^i, \quad i = 0, 1, \dots, m$$

- Assume $p < q$ and $m \approx \infty$

$$\pi_0 = 1 - \rho$$

$$\mathbf{E}[X_n] = \frac{\rho}{1 - \rho} \quad (\text{in steady-state})$$