

Markov Processes – III

Readings: Section 6.4

Lecture outline

- Review of steady-state behavior
- Probability of blocked phone calls
- Calculating absorption probabilities
- Calculating expected time to absorption

- Assume a single class of recurrent states, aperiodic. Then,

$$\lim_{n \rightarrow \infty} r_{ij}(n) = \pi_j$$

where  $\pi_j$  does not depend on the initial conditions

$$\lim_{n \rightarrow \infty} \mathbf{P}(X_n = j | X_0) = \pi_j$$


- $\pi_1, \dots, \pi_m$  can be found as the unique solution of the balance equations

$$\pi_j = \sum_k \pi_k p_{kj}$$

together with

$$\sum_j \pi_j = 1$$

Example




$$\pi_1 = 2/7, \pi_2 = 5/7$$

- Assume process starts at state 1.
- $\mathbf{P}(X_1 = 1, \text{ and } X_{100} = 1) =$
- $\mathbf{P}(X_{100} = 1 \text{ and } X_{101} = 2) =$

The phone company problem

- Calls originate as a Poisson process, rate  $\lambda$ 
  - Each call duration is exponentially distributed (parameter  $\mu$ )
  - $B$  lines available
- Discrete time intervals of (small) length  $\delta$




- Balance equations:  $\lambda \pi_{i-1} = i \mu \pi_i$

$$\pi_i = \pi_0 \frac{\lambda^i}{\mu^i i!} \quad \pi_0 = 1 / \sum_{i=0}^B \frac{\lambda^i}{\mu^i i!}$$

## Calculating absorption probabilities

- What is the probability  $a_i$  that the process eventually settles in state 4, given that the initial state is  $i$ ?




For  $i = 4$ ,  $a_i =$

For  $i = 5$ ,  $a_i =$

$$a_i = \sum_j p_{ij} a_j, \quad \text{for all other } i$$

## Expected time to absorption



- What is the expected number of transitions  $\mu_i$  until the process reaches the absorbing state, given that the initial state is  $i$ ?

$\mu_i = 0$  for  $i =$

For all other  $i$ :  $\mu_i = 1 + \sum_j p_{ij} \mu_j$