

## Lecture 10

Lecturer: Michel X. Goemans

Scribe: Ashish Koul

## 1 Minimum Cost Circulation Problem

**Theorem 1** *Let  $f$  be a circulation. The following are equivalent:*

- (i)  $f$  is of minimum cost.
- (ii)  $G_f$  has no negative cost directed cycles.
- (iii)  $\exists p : c_p(v, w) \geq 0 \quad \forall (v, w) \in E_f$ , where  $c_p(v, w) = c(v, w) + p(v) - p(w)$ .

**Proof:**  $i \Rightarrow ii$  and  $iii \Rightarrow i$  were proven last lecture. All that remains is the proof of  $ii \Rightarrow iii$ :  
 Let  $G'$  be obtained from the residual graph  $G_f$  by adding a vertex  $s$  linked to all other vertices by edges of cost 0 (the costs of these edges do not matter). Let  $p(v)$  be the length of the shortest path from  $s$  to  $v$  in  $G'$  with respect to the costs.  
 These quantities are well-defined since  $G_f$  does not contain any negative cost directed cycles, and every vertex is reachable from  $s$ . By definition of the shortest paths,  $p(w) \leq p(v) + c(v, w) \quad \forall (v, w) \in E_f$ . This implies that  $c_p(v, w) \geq 0 \quad \forall (v, w) \in E_f$ .  $\square$

## 2 Klein's Algorithm for MCCP

**Klein's Cycle canceling algorithm:**

1. Let  $f$  be any circulation.
2. While  $G_f$  contains a negative cycle  $\Gamma$  do  
 push  $\delta = \min_{(v,w) \in \Gamma} u_f(v, w)$  along  $\Gamma$ .

**Argument for Correctness:**

If the algorithm terminates, then the circulation found must be optimum. Furthermore, if all capacities and costs are integers, then the algorithm will terminate.

**Why?**

- $f(v, w)$  is always an integer, thus  $\delta = \min_{(v,w) \in \Gamma} u_f(v, w) \geq 1$
- If  $|c(v, w)| \leq C$  and  $|f(v, w)| \leq U$ , then the absolute value of the cost of the optimal circulation is at most  $mCU$

Therefore, the algorithm terminates after  $O(mCU)$  iterations.

**Remark 1** *If the edge capacities in the graph are irrational, then the algorithm is not correct.*

The cycle canceling algorithm can be applied to the Max-Flow Problem by making appropriate modifications to the graph  $G$ . Let  $G'$  be obtained by setting the cost of all edges within  $G$  to 0. Furthermore, select two vertices  $s$  and  $t$  from within the graph, and add the directed edges  $(s, t)$  and  $(t, s)$ , where  $c(s, t) = 1, c(t, s) = -1$  and both edges have infinite capacity. Now, solving for

the maximum flow between  $s$  to  $t$  is equivalent to solving for the minimum cost circulation, which contains  $s$  and  $t$ . In this circumstance, Klein's Algorithm reduces to the Ford-Fulkerson Algorithm.

**Ford-Fulkerson Augmenting Path Algorithm:**

1. Begin with zero flow:  $f = 0$ .
2. While  $G_f$  contains a directed path  $P$  from  $s$  to  $t$  do  
push  $\delta = \min_{(v,w) \in P} u_f(v, w)$  along  $P$ .

The running time given above for Klein's Cycle-Canceling Algorithm is not polynomial. The negative cost cycle in Klein's Algorithm (or the directed path in the Ford-Fulkerson Algorithm) must be chosen appropriately to insure a polynomial running time.

**Candidates for Cycles in Klein's Algorithm:**

1. The most negative cost cycle in  $G_f$ ?  
*Finding this cycle is an NP-Hard problem, so it would not be a viable choice.*
2. The negative cycle in  $G_f$  which would yield the maximum cost improvement?  
*Finding this cycle is again an NP-Hard problem.*

However for the Max-Flow Problem, this choice reduces to finding the  $st$ -path with maximum residual capacity. Such a path can be found in  $O(m)$  time,  $m = |E|$ . The resulting Max-Flow algorithm is known as the "fattest" path algorithm (Edmonds-Karp '72). The number of iterations necessary is  $O(m \log U)$ , thus the running of the algorithm is  $O(m^2 \log U)$ .

3. Minimum Mean-Cost Cycle?  
Define the mean cost of a cycle  $\Gamma$  to be:

$$\frac{c(\Gamma)}{|\Gamma|} = \frac{\sum_{(v,w) \in \Gamma} c(v, w)}{|\Gamma|} \tag{1}$$

where  $|\Gamma|$  denotes the number of edges in  $\Gamma$ . The minimum mean cost of all cycles of the residual graph  $G_f$  would thus be:

$$\mu(f) = \min_{\text{cycles } \Gamma \text{ in } G_f} \frac{c(\Gamma)}{|\Gamma|} \tag{2}$$

The minimum mean-cost cycle can be determined in strongly polynomial time by using a modified version of the Bellman-Ford Algorithm. More precisely, the minimum mean cost cycle can be found in  $O(mn)$  time. Using this method to solve the Min-Cost Circulation Problem yields the Goldberg-Tarjan Algorithm, which runs in polynomial time. Using this method to solve the Max-Flow Problem yields what is known as the "shortest" augmenting path algorithm (Edmonds-Karp). This Max-Flow Algorithm is able to find the augmenting path in  $O(m)$  time, and requires  $O(mn)$  iterations to arrive at the solution. Thus, the total running time is  $O(m^2n)$ .

### 3 The Goldberg-Tarjan Algorithm

#### Goldberg-Tarjan Algorithm:

1. Begin with zero flow:  $f = 0$ .
2. While  $\mu(f) < 0$  do  
     push  $\delta = \min_{(v,w) \in \Gamma} u_f(v, w)$  along a minimum mean cost cycle  $\Gamma$  of  $G_f$ .

#### Analysis of Goldberg-Tarjan Algorithm:

In order to analyze this algorithm, it is necessary to define the concept of proximity measure for a circulation  $f$ .

**Definition 1** A circulation  $f$  is  $\epsilon$ -optimal if there exists  $p$  such that  $c_p(v, w) \geq -\epsilon \forall (v, w) \in E_f$ .

**Definition 2**  $\epsilon(f) = \text{minimum } \epsilon \text{ such that } f \text{ is } \epsilon\text{-optimal}$ .

The following theorem states that the minimum mean cost  $\mu(f)$  of all cycles in  $G_f$  is equal to  $-\epsilon(f)$ , as defined above.

**Theorem 2** For any circulation  $f$ ,  $\mu(f) = -\epsilon(f)$ .

#### Proof:

- $\mu(f) \geq -\epsilon(f)$   
 By definition, there exists  $p$  such that  $c_p(v, w) \geq -\epsilon(f) \forall (v, w) \in E_f$ . Thus, it is implied that  $c_p(\Gamma) \geq -\epsilon(f)|\Gamma|$  for any directed cycle  $\Gamma \in G_f$ . But for any  $\Gamma \in G_f$ ,  $c(\Gamma) = c_p(\Gamma)$ . Thus, dividing both sides by  $|\Gamma|$  yields that the mean cost of any directed cycle  $\Gamma \in G_f$  is at least  $-\epsilon(f)$ . Therefore,  $\mu(f) \geq -\epsilon(f)$ .
- $\epsilon(f) \leq -\mu(f)$   
 Consider  $\mu(f)$ . For every cycle  $\Gamma \in G_f$ , it is the case that  $\frac{c(\Gamma)}{|\Gamma|} \geq \mu(f)$ . Let  $c'(v, w) = c(v, w) - \mu(f) \forall (v, w) \in E_f$ . With respect to this new cost function  $c'$  every cycle  $\Gamma \in G_f$  will have nonnegative cost. Now, let  $G'$  be obtained by adding a new node  $s$  to  $G_f$  and adding directed edges from  $s$  to  $v \forall v \in V$ , all with zero cost. Let  $p(v)$  be the cost with respect to  $c'$  of the shortest path from  $s$  to  $v$  in the new graph  $G'$ . For all edges  $(v, w)$ ,  $p(w) \leq p(v) + c'(v, w) = p(v) + c(v, w) - \mu(f)$ . This implies that  $c_p(v, w) \geq \mu(f) \forall (v, w) \in E_f$ . Therefore,  $\epsilon(f) \leq -\mu(f)$ .
- $\mu(f) \geq -\epsilon(f)$  and  $\epsilon(f) \leq -\mu(f) \Rightarrow \epsilon(f) = -\mu(f)$ .

□

**Remark 2** Along the minimum mean cost cycle  $\Gamma$ ,  $c_p(v, w) = -\epsilon(f) \forall (v, w) \in \Gamma$ .

Having completed the necessary definitions and proofs, we may now proceed with the analysis of the Goldberg-Tarjan Algorithm. The following theorem considers only one iteration of the algorithm.

**Theorem 3** Let  $f$  be a circulation and let  $f'$  be the circulation obtained by canceling the minimum mean cost cycle  $\Gamma$  of  $G_f$ . Then,  $\epsilon(f') \leq \epsilon(f)$ .

**Proof:** By definition, there exists  $p$  such that  $c_p(v, w) \geq -\epsilon(f) \forall (v, w) \in E_f$ . In the case of the minimum mean cost cycle  $\Gamma$  of  $G_f$ ,  $c_p(v, w) = -\epsilon(f) \forall (v, w) \in \Gamma$ . After performing the one cycle-canceling step, we obtain the new residual graph  $G_{f'}$ . We claim that  $c_p(v, w) \geq -\epsilon(f) \forall (v, w) \in E_{f'}$ . In the case of all edges  $(v, w) \in E_{f'} \cap E_f$ , the claim is certainly true. In the case of all edges

$(v, w) \in E_{f'} \setminus E_f$ , it must be true that  $(w, v) \in \Gamma$ . For all  $(w, v) \in \Gamma$ ,  $c_p(w, v) = -\epsilon(f)$ , and thus  $c_p(v, w) = \epsilon(f) \geq -\epsilon(f)$ . Therefore,  $c_p(v, w) \geq -\epsilon(f)$  holds true for all  $(v, w) \in E_{f'}$ .  $\square$

The above theorem shows that by completing a single iteration of the Goldberg-Tarjan algorithm, it is impossible to generate a new flow which is farther from optimality than the original.