

18.022 Hour Test
October 28, 2005

CLOSED BOOK; NO BOOKS, NOTES, OR CALCULATORS

Name	Rec.Instr.	Rec.Time
------	------------	----------

Please show all your work on this paper, and identify your answers clearly. Use backs of pages if necessary. Points for each question are as shown (for a total of 100 points). If you have difficulty on a problem, go on to the next.

1. (20) Let $f(x,y,z) = (x^2 + y^2 + z^2)^{1/2}$. Use linear approximation to estimate $f(Q) - f(P)$, the increase in the value of f from $P = (2,2,1)$ to $Q = (2.1, 1.9, 0.8)$.

SCORE

1.

2.

3.

4.

5.

6.

TOTAL:

2. (10) The temperature at each point (x,y,z) in space at time t is given by the function $T(x,y,z,t)$. The path of a moving particle is given by the position vector $\vec{R}(t) = x(t)\hat{i} + y(t)\hat{j} + z(t)\hat{k}$. Use the chain rule to give an expression, in terms of derivatives of the functions T , x , y , and z , for the instantaneous rate per unit time at which the temperature experienced by the particle is increasing.

3. (10) Let the equations $f(x,y,z) = 0$ and $x = g(y)$, where $f(x,y,z)$ and $g(y)$ are given by mathematical formulas, define y implicitly as a function of x and z . Find $\partial y / \partial x$ in terms of the formal derivatives f_x , f_y , f_z , and g_y . (Suggestion: use the elimination method.)

4. (20) Consider the scalar field given by $f(x,y) = xy^2$ in the xy plane. If we limit our attention to points on the circle $x^2 + y^2 = 9$, then f has six constrained critical points. Use a Lagrange multiplier to find these points and show them in a figure. Indicate on the figure which points are global maximum points, global minimum points, local maximum but not global maximum points, and local minimum but not global minimum points. (Hint: use the max-min existence theorem.)

5. (20) A solid rectangular block of constant density has total mass M , length = a , width = b , and height = c . Find its moment of inertia about the vertical axis through the centers of the two opposite faces of length a and width b . Express your answer in terms of M , a , b , and (if necessary) c . (The integration is easiest in Cartesian coordinates. Use the z axis as the axis of the moment of inertia.)

6. (20) You are given the double integral $\iint_R \frac{y-x}{y+x} dA$ where R is the triangular region in

the xy plane with vertices $(0, 0)$, $(1/2, 1/2)$, and $(0, 1)$.

(a) (15 points) Express this integral as an iterated integral in new variables u and v such that $u = y + x$ and $v = y - x$. Be sure to show limits of integration.

(b) (5 points) Evaluate the expression found in (a).