

# **Lecture 2**

## **Orthogonal Vectors and Matrices, Norms**

MIT 18.335J / 6.337J

Introduction to Numerical Methods

Per-Olof Persson

September 12, 2006

# Transpose and Adjoint

- For real  $A$ , the *transpose* of  $A$  is obtained by interchanging rows/columns

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \implies A^T = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{12} & a_{22} & a_{32} \end{bmatrix}$$

- The *adjoint* or *hermitian conjugate* also takes complex conjugate

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix} \implies A^* = \begin{bmatrix} \bar{a}_{11} & \bar{a}_{21} & \bar{a}_{31} \\ \bar{a}_{12} & \bar{a}_{22} & \bar{a}_{32} \end{bmatrix}$$

- If real  $A = A^T$ , then  $A$  is *symmetric*. If  $A = A^*$ , then  $A$  is *hermitian*.

# Inner Product

- *Inner product* of two column vectors  $x, y \in \mathbb{C}^m$

$$x^* y = \sum_{i=1}^m \bar{x}_i y_i$$

- *Euclidean length* of  $x$

$$\|x\| = \sqrt{x^* x} = \left( \sum_{i=1}^m |x_i|^2 \right)^{1/2}$$

- *Angle*  $\alpha$  between  $x, y$

$$\cos \alpha = \frac{x^* y}{\|x\| \|y\|}$$

# In MATLAB

Quantity	MATLAB Syntax	Comment
Transpose $A^T$	<code>A . '</code>	Transpose only
Adjoint $A^*$	<code>A '</code>	Transpose + complex conjugate
Inner product $x^*y$	<code>x' * y</code> or <code>dot(x, y)</code>	' * assumes column vectors
Length $\ x\ $	<code>sqrt(x' * x)</code> or <code>norm(x)</code>	' * assumes column vector

# Orthogonal Vectors

- The vectors  $x, y \in \mathbb{R}^m$  are *orthogonal* if

$$x^*y = 0$$

- The sets of vectors  $X, Y$  are orthogonal if

every  $x \in X$  is orthogonal to every  $y \in Y$

- A set of (nonzero) vectors  $S$  is orthogonal if

vectors pairwise orthogonal, i.e., for  $x, y \in S, x \neq y \Rightarrow x^*y = 0$

and *orthonormal* if, in addition,

every  $x \in S$  has  $\|x\| = 1$

# Orthogonal and Unitary Matrices

- A square matrix  $Q \in \mathbb{C}^{m \times m}$  is *unitary* (*orthogonal* in real case), if

$$Q^* = Q^{-1}$$

- For unitary  $Q$

$$Q^* Q = I, \text{ or } q_i^* q_j = \delta_{ij}$$

- Interpretation of unitary-times-vector product:

$$x = Q^* b = \text{solution to } Qx = b$$

= the vector of coefficients of the expansion of  $b$

in the basis of columns of  $Q$


# Preservation of Geometry Structure

- Inner product is preserved under multiplication by unitary  $Q$


$$(Qx)^*(Qy) = x^*Q^*Qy = x^*y$$

- Therefore, *lengths of vectors* and *angles between vectors* are preserved
- A real orthogonal  $Q$  is either a rigid rotation or reflection

**Rotation**



**Reflection**



# Vector Norms

- A *norm* is a function  $\| \cdot \| : \mathbb{C}^m \rightarrow \mathbb{R}$  satisfying

$$(1) \quad \|x\| \geq 0, \text{ and } \|x\| = 0 \text{ only if } x = 0$$

$$(2) \quad \|x + y\| \leq \|x\| + \|y\|$$

$$(3) \quad \|\alpha x\| = |\alpha| \|x\|$$

- Example: The Euclidean length function

$$\|x\|_2 = \sqrt{x^* x}$$

- $\| \cdot \|_2$  is a special case of the  $p$ -norms

$$\|x\|_p = \left( \sum_{i=1}^m |x_i|^p \right)^{1/p} \quad (1 \leq p < \infty)$$

# Examples of Vector Norms

$$\|x\|_1 = \sum_{i=1}^m |x_i|$$

$$\|x\|_2 = \left( \sum_{i=1}^m |x_i|^2 \right)^{1/2} = \sqrt{x^* x}$$

$$\|x\|_\infty = \max_{1 \leq i \leq m} |x_i|$$

$$\|x\|_W = \|Wx\|_2 = \left( \sum_{i=1}^m |w_i x_i|^2 \right)^{1/2}$$

# Verification of Norm Conditions

- Show that  $\|x\|_1 = \sum_{i=1}^m |x_i|$  is a norm

(1)

$$\|x\|_1 = |x_1| + |x_2| + \cdots + |x_m| \geq 0, \text{ equality only if } x = 0$$

(2)

$$\begin{aligned} \|x + y\|_1 &= \sum_{i=1}^m |x_i + y_i| \leq \sum_{i=1}^m (|x_i| + |y_i|) \\ &= \sum_{i=1}^m |x_i| + \sum_{i=1}^m |y_i| = \|x\|_1 + \|y\|_1 \end{aligned}$$

(3)

$$\|\alpha x\|_1 = \sum_{i=1}^m |\alpha x_i| = \sum_{i=1}^m |\alpha| |x_i| = |\alpha| \sum_{i=1}^m |x_i| = |\alpha| \|x\|_1$$


# Induced Matrix Norms

- For a matrix  $A \in \mathbb{C}^{m \times n}$ , the *induced matrix norm* is

$$\|A\|_{(m,n)} = \sup_{\substack{x \in \mathbb{C}^n \\ x \neq 0}} \frac{\|Ax\|_{(m)}}{\|x\|_{(n)}} = \sup_{\substack{x \in \mathbb{C}^n \\ \|x\|_{(n)}=1}} \|Ax\|_{(m)}$$

where  $\|\cdot\|_{(n)}$  and  $\|\cdot\|_{(m)}$  are given vector norms

- The “maximum stretching” by  $A$



# Examples of Matrix Norms

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^m |a_{ij}|$$

“maximum column sum”

$$\|A\|_\infty = \max_{1 \leq i \leq m} \sum_{j=1}^n |a_{ij}|$$

“maximum row sum”

$$\|A\|_F = \left( \sum_{i=1}^m \sum_{j=1}^n |a_{ij}|^2 \right)^{1/2}$$

The Frobenius norm

$$\|A\|_2 = \sqrt{\lambda_{\max}(A^*A)}$$

More later

# Properties of Matrix Norms

- *Bound on Matrix Product* - Induced norms and Frobenius norm satisfy

$$\|AB\| \leq \|A\|\|B\|$$

but some matrix norms do not!

- *Invariance under Unitary Multiplication* - For  $A \in \mathbb{C}^{m \times n}$  and unitary  $Q \in \mathbb{C}^{m \times m}$ , we have

$$\|QA\|_2 = \|A\|_2, \quad \|QA\|_F = \|A\|_F$$

*Proof.* Since  $\|Qx\|_2 = \|x\|_2$  (inner product is preserved), the first result follows from the definition of induced norm. For the Frobenius norm,

$$\begin{aligned} \|QA\|_F &= \sqrt{\text{tr}((QA)^*(QA))} = \sqrt{\text{tr}(A^*Q^*QA)} \\ &= \sqrt{\text{tr}(A^*A)} = \|A\|_F \end{aligned}$$

# In MATLAB

Quantity	MATLAB Syntax
$\ x\ _1$	<code>sum(abs(x))</code> or <code>norm(x,1)</code>
$\ x\ _2$	<code>sqrt(x'*x)</code> or <code>norm(x)</code>
$\ x\ _p$	<code>sum(abs(x).^p).^(1/p)</code> or <code>norm(x,p)</code>
$\ x\ _\infty$	<code>max(abs(x))</code> or <code>norm(x,inf)</code>
$\ A\ _1$	<code>max(sum(abs(A),1))</code> or <code>norm(A,1)</code>
$\ A\ _2$	<code>norm(A)</code>
$\ A\ _\infty$	<code>max(sum(abs(A),2))</code> or <code>norm(A,inf)</code>
$\ A\ _F$	<code>sqrt(A(:)'*A(:))</code> or <code>norm(A,'fro')</code>