

Lecture 23

Arnoldi/Lanczos Iterations

MIT 18.335J / 6.337J

Introduction to Numerical Methods

Per-Olof Persson

December 5, 2006

The Arnoldi Iteration

- The Arnoldi process reduces a general, nonsymmetric A to Hessenberg form by similarity transforms: $A = QHQ^*$
- Allows for reduced factorizations by a Gram-Schmidt-style iteration instead of Householder reflections
- Let Q_n be the $m \times n$ matrix with the first n columns of Q , and consider the first n columns of $AQ = QH$, or $AQ_n = Q_{n+1}\tilde{H}_n$:

$$\begin{bmatrix} A \end{bmatrix} \begin{bmatrix} q_1 & \cdots & q_n \end{bmatrix} = \begin{bmatrix} q_1 & \cdots & q_{n+1} \end{bmatrix} \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ h_{21} & & \\ & \ddots & \\ & & h_{n+1,n} \end{bmatrix}$$

The Arnoldi Algorithm

- The n th column of $AQ_n = Q_{n+1}\tilde{H}_n$ gives

$$Aq_n = h_{1n}q_1 + \cdots + h_{nn}q_n + h_{n+1,n}q_{n+1}$$

which can be used to compute q_{n+1} similarly to modified Gram-Schmidt:

Algorithm: Arnoldi Iteration

$$b = \text{arbitrary}, q_1 = b/\|b\|$$

for $n = 1, 2, 3, \dots$

$$v = Aq_n$$

for $j = 1$ **to** n

$$h_{jn} = q_j^* v$$

$$v = v - h_{jn}q_j$$

$$h_{n+1,n} = \|v\|$$

$$q_{n+1} = v/h_{n+1,n}$$

QR Factorization of Krylov Matrix

- The vectors q_j from Arnoldi are orthonormal bases of the successive *Krylov subspaces*:

$$\mathcal{K}_n = \langle b, Ab, \dots, A^{n-1}b \rangle = \langle q_1, q_2, \dots, q_n \rangle \subseteq \mathbb{C}^m$$

- Q_n is the reduced QR factorization $K_n = Q_n R_n$ of the *Krylov matrix*:

$$K_n = \left[\begin{array}{c|c|c|c} b & Ab & \dots & A^{n-1}b \end{array} \right]$$

- The projection of A onto this space gives $n \times n$ Hessenberg matrix $H_n = Q_n^* A Q_n$, whose eigenvalues may be good approximations of A 's

Symmetric Matrices and the Lanczos Iteration

- For symmetric A , H_n reduces to tridiagonal T_n , and q_{n+1} can be computed by a three-term recurrence:

$$Aq_n = \beta_{n-1}q_{n-1} + \alpha_n q_n + \beta_n q_{n+1}$$

Algorithm: Lanczos Iteration

$$\beta_0 = 0, q_0 = 0, b = \text{arbitrary}, q_1 = b/\|b\|$$

for $n = 1, 2, 3, \dots$

$$v = Aq_n$$

$$\alpha_n = q_n^T v$$

$$v = v - \beta_{n-1}q_{n-1} - \alpha_n q_n$$

$$\beta_n = \|v\|$$

$$q_{n+1} = v/\beta_n$$