

Review Paramagnet: a collection of non-interacting magnetic moments

Quantum Mechanics


$$\vec{\mu} = g_J \mu_B \vec{J}$$

$\Rightarrow 2J + 1$  levels

The topic was saved until the last because it is subtle.

First  $J = 1/2$ , Microcanonical:  $S(M) \rightarrow M(T, H)$

Second  $J = 1/2$ , Canonical:  $p(m_J) \rightarrow \langle \mu \rangle \rightarrow M(T, H)$



Third  $J > 1/2$ , Canonical:  $\langle \mu \rangle \rightarrow M(T, H)$

$$Z = Z(\eta = g\mu_B H/kT)$$

So what's the problem?

$$\langle \mathcal{H} \rangle^{\text{S. M.}} \neq U^{\text{Thermo}}$$

$$= U^{\text{assembly}} + (-\vec{H} \cdot \vec{M})^{\text{position}} \quad (\text{for } \delta W = H dM)$$


$$\langle \mathcal{H} \rangle = -\frac{1}{Z} \left( \frac{\partial Z}{\partial \beta} \right)_H = -\frac{1}{Z} \frac{dZ}{d\eta} \underbrace{\left( \frac{\partial \eta}{\partial \beta} \right)_H}_{g\mu_B H} = -HM \quad \checkmark$$

$\underbrace{Z}_{M/g\mu_B}$

## Entropy of a Quantum Paramagnet

- When is  $-kT \ln Z \neq F$  ?
- How is a paramagnet like a sponge?

# HIGH AND LOW TEMPERATURE BEHAVIOR OF A QUANTUM PARAMAGNET



ENERGY LEVELS ALMOST  
EQUALLY POPULATED,  
CURIE LAW BEHAVIOR

MOMENT ALMOST SATURATED,  
ENERGY GAP BEHAVIOR

$$S(kT \ll g\mu_B H) \rightarrow Nk \ln(1) = 0$$

$$S(kT \gg g\mu_B H) \rightarrow Nk \ln(2J + 1)$$

$$Z_1 = \sum_{m=-J}^J (e^\eta)^m \quad \eta \equiv g\mu_B H/kT$$

Try

$$-kT \ln Z = F = \underbrace{U}_0 - TS \Rightarrow S = k \ln Z = Nk \ln Z_1$$

$$S(kT \ll g\mu_B H) \rightarrow Nk \ln(1) = 0$$

$$S(kT \gg g\mu_B H) \rightarrow Nk \ln(2J + 1) \quad \underline{Nk \ln(2J + 1) \text{ O.K.}}$$

$$Z_1 = \sum_{m=-J}^J (e^\eta)^m \quad \eta \equiv g\mu_B H/kT$$

Try

$$-kT \ln Z = F = \underbrace{U}_0 - TS \Rightarrow S = k \ln Z = Nk \ln Z_1$$

$$S(kT \ll g\mu_B H) \rightarrow Nk \ln(1) = 0 \quad \underline{NkJ(g\mu_B H/kT) \text{ wrong!}}$$

$$S(kT \gg g\mu_B H) \rightarrow Nk \ln(2J + 1) \quad \underline{Nk \ln(2J + 1) \text{ O.K.}}$$

$$Z_1 = \sum_{m=-J}^J (e^\eta)^m \quad \eta \equiv g\mu_B H/kT$$

Try

$$-kT \ln Z \underbrace{\equiv F}_{\text{wrong}} = \underbrace{U}_0 - TS \Rightarrow S = k \ln Z = \underbrace{Nk \ln Z_1}_{\text{wrong}}$$

In the derivation of the canonical ensemble we found



$$-kT \ln Z = \langle E_1 \rangle - TS_1 \text{ where } \langle E_1 \rangle = \langle \mathcal{H}(\{p, q\}) \rangle$$

Then we set  $\langle E_1 \rangle = U$ . But in the paramagnet  $\langle E_1 \rangle = U - HM$ , thus

$$-kT \ln Z = U - HM - TS = G(T, H) \text{ for our model.}$$

$$\Rightarrow S = k \ln Z - HM/T = \underline{Nk \ln Z_1(\eta) - NkJ \eta B_J(\eta)}$$

# ADIABATIC DEMAGNETIZATION (MAGNETIC COOLING)




$$H = H_{\text{initial}}$$

$$\left. \begin{array}{l} S_M \sim 0 \\ S_S \text{ high} \end{array} \right\} S_{\text{total}}$$

$$T_S = T_{\text{initial}}$$

# ADIABATIC DEMAGNETIZATION (MAGNETIC COOLING)

INITIAL COOLING  
STAGE  
 $T_{\text{initial}}$



$$2J+1 \text{ --- } \otimes \text{ ---}$$

$$kT \gg g\mu_B H$$

$$H \sim 0$$

$$S_M \sim Nk \ln(2J+1) \left. \vphantom{S_M} \right\} S_{\text{total}}$$

$$S_S \text{ low}$$

$$T_S \ll T_{\text{initial}}$$

## Electronic Example, CMN

Cerium Manganese Nitrate



$\text{Ce}^{+++}$        $J=5/2$        $T_{\text{ordering}} \sim 1.9 \text{ mK}$

Cools  $^3\text{He}$  and samples therein to  $\sim 2 \text{ mK}$ .

## Nuclear Example, Cu

Metallic Copper

Cu       $I=3/2$        $T_{\text{ordering}} \sim 1 \mu\text{K}$

Cools Cu electrons and lattice to  $\sim 10 \mu\text{K}$ .