

Practice Exam 1: Long List

18.05, Spring 2022

This is a big list of practice problems for Exam 1. It includes all the problems in other sets of practice problems and many more!

1 Counting and Probability

Problem 1. A full house in poker is a hand where three cards share one rank and two cards share another rank. How many ways are there to get a full-house? What is the probability of getting a full-house?

Problem 2. There are 3 arrangements of the word DAD, namely DAD, ADD, and DDA. How many arrangements are there of the word PROBABILITY?

Problem 3. (a) How many ways can you arrange the letters in the word STATISTICS? (e.g. SSSTTTIIAC counts a one arrangement.)

(b) If all arrangements are equally likely, what is the probability the two 'i's are next to each other.

Problem 4. In a ballroom dancing class the students are divided into group A and group B . There are six people in group A and seven in group B . If four A s and four B s are chosen and paired off, how many pairings are possible?

Problem 5. Suppose you pick two cards from a deck of 52 playing cards. What is the probability that they are both queens?

Problem 6. Suppose that there are ten students in a classroom. What is the probability that no two of them have a birthday in the same month?

Problem 7. 20 politicians are having a tea party, 6 Democrats and 14 Republicans. To prepare, they need to choose:

3 people to set the table, 2 people to boil the water, 6 people to make the scones. Each person can only do 1 task. (Note that this doesn't add up to 20. The rest of the people don't help.)

(a) In how many different ways can they choose which people perform these tasks?

(b) Suppose that the Democrats all hate tea. If they only give tea to 10 of the 20 people, what is the probability that they only give tea to Republicans?

(c) If they only give tea to 10 of the 20 people, what is the probability that they give tea to 9 Republicans and 1 Democrat?

Problem 8. Let A and B be two events. Suppose the probability that neither A or B occurs is $2/3$. What is the probability that one or both occur?

Problem 9. Let C and D be two events with $P(C) = 0.25$, $P(D) = 0.45$, and $P(C \cap D) = 0.1$. What is $P(C^c \cap D)$?

Problem 10. You roll a four-sided die 3 times. For this problem we'll use the sample space with 64 equally likely outcomes.

- (a) Write down this sample space in set notation.
- (b) List all the outcomes in each of the following events.
 - (i) $A =$ 'Exactly 2 of the 3 rolls are fours'
 - (ii) $B =$ 'At least 2 of the 3 rolls are fours'
 - (iii) $C =$ 'Exactly 1 of the second and third rolls is a 4'
 - (iv) $A \cap C$

Problem 11. Suppose we have 8 teams labeled T_1, \dots, T_8 . Suppose they are ordered by placing their names in a hat and drawing the names out one at a time.

- (a) How many ways can it happen that all the odd numbered teams are in the odd numbered slots and all the even numbered teams are in the even numbered slots?
- (b) What is the probability of this happening?

2 Conditional Probability and Bayes' Theorem

Problem 12. More cards! Suppose you want to divide a 52 card deck into four hands with 13 cards each. What is the probability that each hand has a king?

Problem 13. Suppose you are taking a multiple-choice test with c choices for each question. In answering a question on this test, the probability that you know the answer is p . If you don't know the answer, you choose one at random. What is the probability that you knew the answer to a question, given that you answered it correctly?

Problem 14. Corrupted by their power, the judges running the popular game show *America's Next Top Mathematician* have been taking bribes from many of the contestants. Each episode, a given contestant is either allowed to stay on the show or is kicked off.

If the contestant has been bribing the judges they will be allowed to stay with probability 1. If the contestant has not been bribing the judges, they will be allowed to stay with probability $1/3$.

Suppose that $1/4$ of the contestants have been bribing the judges. The same contestants bribe the judges in both rounds, i.e., if a contestant bribes them in the first round, they bribe them in the second round too (and vice versa).

- (a) If you pick a random contestant who was allowed to stay during the first episode, what is the probability that they were bribing the judges?
- (b) If you pick a random contestant, what is the probability that they are allowed to stay during both of the first two episodes?

(c) If you pick random contestant who was allowed to stay during the first episode, what is the probability that they get kicked off during the second episode?

Problem 15. Consider the Monty Hall problem. Let's label the door with the car behind it a and the other two doors b and c . In the game the contestant chooses a door and then Monty chooses a door, so we can label each outcome as 'contestant followed by Monty', e.g ab means the contestant chose a and Monty chose b .

- (a) Make a 3×3 probability table showing probabilities for all possible outcomes.
- (b) Make a probability tree showing all possible outcomes.
- (c) Suppose the contestant's strategy is to switch. List all the outcomes in the event 'the contestant wins a car'. What is the probability the contestant wins?
- (d) Redo part (c) with the strategy of not switching.

Problem 16. Two dice are rolled.

A = 'sum of two dice equals 3'

B = 'sum of two dice equals 7'

C = 'at least one of the dice shows a 1'

- (a) What is $P(A|C)$?
- (b) What is $P(B|C)$?
- (c) Are A and C independent? What about B and C ?

Problem 17. There is a screening test for prostate cancer that looks at the level of PSA (prostate-specific antigen) in the blood. There are a number of reasons besides prostate cancer that a man can have elevated PSA levels. In addition, many types of prostate cancer develop so slowly that that they are never a problem. Unfortunately there is currently no test to distinguish the different types and using the test is controversial because it is hard to quantify the accuracy rates and the harm done by false positives.

For this problem we'll call a positive test a true positive if it catches a dangerous type of prostate cancer. We'll assume the following numbers:

Rate of prostate cancer among men over 50 = 0.0005

True positive rate for the test = 0.9

False positive rate for the test = 0.01

Let T be the event a man has a positive test and let D be the event a man has a dangerous type of the disease. Find $P(D|T)$ and $P(D|T^c)$.

Problem 18. A multiple choice exam has 4 choices for each question. A student has studied enough so that the probability they will know the answer to a question is 0.5, the probability that they will be able to eliminate one choice is 0.25, otherwise all 4 choices seem equally plausible. If they know the answer they will get the question right. If not they have to guess from the 3 or 4 choices.

As the teacher you want the test to measure what the student knows. If the student answers a question correctly what's the probability they knew the answer?

Problem 19. Suppose you have an urn containing 7 red and 3 blue balls. You draw three balls at random. On each draw, if the ball is red you set it aside and if the ball is blue you put it back in the urn. What is the probability that the third draw is blue?

(If you get a blue ball it counts as a draw even though you put it back in the urn.)

Problem 20. Some games, like tennis or ping pong, reach a state called *deuce*. This means that the score is tied and a player wins the game when they get *two* points ahead of the other player. Suppose the probability that you win a point is p and this is true independently for all points. If the game is at deuce what is the probability you win the game?

This is a tricky problem, but amusing if you like puzzles.

Problem 21. (Bayes formula)

A student takes a multiple-choice exam. Suppose for each question they either know the answer or gamble and choose an option at random. Further suppose that if they know the answer, the probability of a correct answer is 1, and if they gamble, this probability is $1/4$. To pass, students need to answer at least 60% of the questions correctly. The student has “studied for a minimal pass,” i.e., with probability 0.6 they know the answer to a question. For a single question, given that they answer it correctly, what is the probability that they actually *knew* the answer?

3 Independence

Problem 22. Suppose that $P(A) = 0.4$, $P(B) = 0.3$ and $P((A \cup B)^C) = 0.42$. Are A and B independent?

Problem 23. Suppose now that events A, B and C are *mutually independent* with

$$P(A) = 0.3, \quad P(B) = 0.4, \quad P(C) = 0.5.$$

Compute the following: (Hint: Use a Venn diagram)

(i) $P(A \cap B \cap C^c)$ (ii) $P(A \cap B^c \cap C)$ (iii) $P(A^c \cap B \cap C)$

Problem 24. You roll a twenty-sided die. Determine whether the following pairs of events are independent.

- (a) ‘You roll an even number’ and ‘You roll a number less than or equal to 10’.
- (b) ‘You roll an even number’ and ‘You roll a prime number’.

Problem 25. Suppose A and B are events with $0 < P(A) < 1$ and $0 < P(B) < 1$.

- (a) If A and B are disjoint can they be independent?
- (b) If A and B are independent can they be disjoint?
- (c) If $A \subset B$ can they be independent?

4 Expectation and Variance

Problem 26. Directly from the definitions of expected value and variance, compute $E[X]$ and $\text{Var}(X)$ when X has probability mass function given by the following table:

X	-2	-1	0	1	2
pmf	1/15	2/15	3/15	4/15	5/15

Problem 27. Suppose that X takes values between 0 and 1 and has probability density function $2x$. Compute $\text{Var}(X)$ and $\text{Var}(X^2)$.

Problem 28. The random variable X takes values -1, 0, 1 with probabilities 1/8, 2/8, 5/8 respectively.

- Compute $E[X]$.
- Give the pmf of $Y = X^2$ and use it to compute $E[Y]$.
- Instead, compute $E[X^2]$ directly from an extended table.
- Compute $\text{Var}(X)$.

Problem 29. Suppose X is a random variable with $E[X] = 5$ and $\text{Var}(X) = 2$. What is $E[X^2]$?

Problem 30. Compute the expectation and variance of a Bernoulli(p) random variable.

Problem 31. Suppose 100 people all toss a hat into a box and then proceed to randomly pick out of a hat. What is the expected number of people to get their own hat back.

Hint: express the number of people who get their own hat as a sum of random variables whose expected value is easy to compute.

Problem 32. Suppose I play a gambling game with even odds. So, I can wager b dollars and I either win or lose b dollars with probability $p = 0.5$.

I employ the following strategy to try to guarantee that I win some money.

I bet \$1; if I lose, I double my bet to \$2, if I lose I double my bet again. I continue until I win. Eventually I'm sure to win a bet and net \$1 (run through the first few rounds and you'll see why this is the net).

If this really worked casinos would be out of business. Our goal in this problem is to understand the flaw in the strategy.

- Let X be the amount of money bet on the last game (the one I win). X takes values 1, 2, 4, 8, Determine the probability mass function for X . That is, find $p(2^k)$, where k is in $\{0, 1, 2, \dots\}$.
- Compute $E[X]$.
- Use your answer in part (b) to explain why the strategy is a bad one.

Problem 33. Suppose you roll a fair 6-sided die 100 times (independently), and you get \$3 every time you roll a 6.

Let X_1 be the number of dollars you win on rolls 1 through 25.

Let X_2 be the number of dollars you win on rolls 26 through 50.

Let X_3 be the number of dollars you win on rolls 51 through 75.

Let X_4 be the number of dollars you win on rolls 76 through 100.

Let $X = X_1 + X_2 + X_3 + X_4$ be the total number of dollars you win over all 100 rolls.

- (a) What is the probability mass function of X ?
- (b) What is the expectation and variance of X ?
- (c) Let $Y = 4X_1$. (So instead of rolling 100 times, you just roll 25 times and multiply your winnings by 4.)
 - (i) What are the expectation and variance of Y ?
 - (ii) How do the expectation and variance of Y compare to those of X ? (That is, are they bigger, smaller, or equal?) Explain (briefly) why this makes sense.

5 Probability Mass Functions, Probability Density Functions and Cumulative Distribution Functions

Problem 34. Suppose that $X \sim \text{Bin}(n, 0.5)$. Find the probability mass function of $Y = 2X$.

- Problem 35.** (a) Suppose that X is uniform on $[0, 1]$. Compute the pdf and cdf of X .
 (b) If $Y = 2X + 5$, compute the pdf and cdf of Y .

Problem 36. (a) Suppose that X has probability density function $f_X(x) = \lambda e^{-\lambda x}$ for $x \geq 0$. Compute the cdf, $F_X(x)$.

- (b) If $Y = X^2$, compute the pdf and cdf of Y .

Problem 37. Suppose that X is a random variable that takes on values 0, 2 and 3 with probabilities 0.3, 0.1, 0.6 respectively. Let $Y = 3(X - 1)^2$.

- (a) What is the expectation of X ?
- (b) What is the variance of X ?
- (c) What is the expectation of Y ?
- (d) Let $F_Y(t)$ be the cumulative density function of Y . What is $F_Y(7)$?

Problem 38. Let T be the waiting time for customers in a queue. Suppose that T is exponential with pdf $f(t) = 2e^{-2t}$ on $[0, \infty)$.

Find the pdf of the rate at which customers are served $R = 1/T$.

Problem 39. A continuous random variable X has PDF $f(x) = x + ax^2$ on $[0, 1]$

Find a , the CDF and $P(0.5 < X < 1)$.

Problem 40. (PMF of a sum)

Suppose X and Y are independent and $X \sim \text{Bernoulli}(1/2)$ and $Y \sim \text{Bernoulli}(1/3)$. Determine the pmf of $X + Y$

Problem 41. Let X be a discrete random variable with pmf p given by:

x	-2	-1	0	1	2
$p(x)$	1/15	2/15	3/15	4/15	5/15


- (a) Let $Y = X^2$. Find the pmf of Y .
 (b) Find the value the cdf of X at $-3/2, 3/4, 7/8, 1, 1.5, 5$.
 (c) Find the value the cdf of Y at $-3/2, 3/4, 7/8, 1, 1.5, 5$.

Problem 42. Suppose that the cdf of X is given by:

$$F(a) = \begin{cases} 0 & \text{for } a < 0 \\ \frac{1}{5} & \text{for } 0 \leq a < 2 \\ \frac{2}{5} & \text{for } 2 \leq a < 4 \\ 1 & \text{for } a \geq 4. \end{cases}$$

Determine the pmf of X .

Problem 43. For each of the following say whether it can be the graph of a cdf. If it can be, say whether the variable is discrete or continuous.



Problem 44. Suppose X has range $[0,1]$ and has cdf

$$F(x) = x^2 \quad \text{for } 0 \leq x \leq 1.$$

Compute $P(\frac{1}{2} < X < \frac{3}{4})$.

Problem 45. Let X be a random variable with range $[0, 1]$ and cdf

$$F(X) = 2x^2 - x^4 \quad \text{for } 0 \leq x \leq 1.$$

- (a) Compute $P(\frac{1}{4} \leq X \leq \frac{3}{4})$.
 (b) What is the pdf of X ?

6 Distributions with Names

Problem 46. Exponential Distribution

Suppose that buses arrive are scheduled to arrive at a bus stop at noon but are always X minutes late, where X is an exponential random variable with probability density function $f_X(x) = \lambda e^{-\lambda x}$. Suppose that you arrive at the bus stop precisely at noon.

- (a) Compute the probability that you have to wait for more than five minutes for the bus to arrive.
 (b) Suppose that you have already waiting for 10 minutes. Compute the probability that you have to wait an additional five minutes or more.

Problem 47. Normal Distribution: Throughout these problems, let ϕ and Φ be the pdf and cdf, respectively, of the standard normal distribution. Suppose Z is a standard normal random variable and let $X = 3Z + 1$.

- (a) Express $P(X \leq x)$ in terms of Φ
 (b) Differentiate the expression from (a) with respect to x to get the pdf of X , $f(x)$. Remember that $\Phi'(z) = \phi(z)$ and don't forget the chain rule
 (c) Find $P(-1 \leq X \leq 1)$
 (d) Recall that the probability that Z is within one standard deviation of its mean is approximately 68%. What is the probability that X is within one standard deviation of its mean?

Problem 48. Transforming Normal Distributions

Suppose $Z \sim N(0,1)$ and $Y = e^Z$.

- (a) Find the cdf $F_Y(a)$ and pdf $f_Y(y)$ for Y . (For the CDF, the best you can do is write it in terms of Φ the standard normal cdf.)
 (b) We don't have a formula for $\Phi(z)$ so we don't have a formula for quantiles. So we have to write quantiles *in terms* of Φ^{-1} .
 (i) Write the 0.33 quantile of Z in terms of Φ^{-1}
 (ii) Write the 0.9 quantile of Y in terms of Φ^{-1} .
 (iii) Find the median of Y .

Problem 49. (Random variables derived from normal random variables)

Let X_1, X_2, \dots, X_n be i.i.d. $N(0, 1)$ random variables.

Let $Y_n = X_1^2 + \dots + X_n^2$.

(a) Use the formula $\text{Var}(X_j) = E[X_j^2] - E[X_j]^2$ to show $E[X_j^2] = 1$.

(b) Set up an integral in x for computing $E[X_j^4]$.

For 3 extra credit points, use integration by parts show $E[X_j^4] = 3$.

(If you don't do this, you can still use this result in part c.)

(c) Deduce from parts (a) and (b) that $\text{Var}(X_j^2) = 2$.

(d) Use the Central Limit Theorem to approximate $P(Y_{100} > 110)$.

Problem 50. More Transforming Normal Distributions

(a) Suppose Z is a standard normal random variable and let $Y = aZ + b$, where $a > 0$ and b are constants.

Show $Y \sim N(b, a^2)$ (remember our notation for normal distributions uses mean and variance).

(b) Suppose $Y \sim N(\mu, \sigma^2)$. Show $\frac{Y - \mu}{\sigma}$ follows a standard normal distribution.

Problem 51. (Sums of normal random variables)

Let X, Y be independent random variables where $X \sim N(2, 5)$ and $Y \sim N(5, 9)$ (we use the notation $N(\mu, \sigma^2)$). Let $W = 3X - 2Y + 1$.

(a) Compute $E[W]$ and $\text{Var}(W)$.

(b) It is known that the sum of independent normal distributions is normal. Compute $P(W \leq 6)$.

Problem 52. Let $X \sim U(a, b)$. Compute $E[X]$ and $\text{Var}(X)$.

Problem 53. In $n + m$ independent Bernoulli(p) trials, let S_n be the number of successes in the first n trials and T_m the number of successes in the last m trials.

(a) What is the distribution of S_n ? Why?

(b) What is the distribution of T_m ? Why?

(c) What is the distribution of $S_n + T_m$? Why?

(d) Are S_n and T_m independent? Why?

Problem 54. Compute the median for the exponential distribution with parameter λ .

Problem 55. Pareto and the 80-20 rule.

Pareto was an economist who used the Pareto distribution to model the wealth in a society. For a fixed baseline m , the Pareto density with parameter α is

$$f(x) = \frac{\alpha m^\alpha}{x^{\alpha+1}} \quad \text{for } x \geq m.$$

Assume X is a random variable that follows such a distribution.

- (a) Compute $P(X > a)$ (you may assume $a \geq m$).
- (b) Pareto's principle is often paraphrased as the 80-20 rule. That is, 80% of the wealth is owned by 20% of the people. The rule is only exact for a Pareto distribution with $\alpha = \log(5)/\log(4) = 1.16$.

Suppose $\alpha = m = 1$. Compute the 0.80 quantile for the Pareto distribution.

In general, many phenomena follow the power law described by $f(x)$. You can look up 'Pareto principle' in Wikipedia to read more about this.

7 Joint Probability, Covariance, Correlation

Problem 56. (Another Arithmetic Puzzle)

Let X and Y be two independent Bernoulli(0.5) random variables. Define S and T by:

$$S = X + Y \quad \text{and} \quad T = X - Y.$$

- (a) Find the joint and marginal pmf's for S and T .
- (b) Are S and T independent.

Problem 57. Data is taken on the height and shoe size of a sample of MIT students. Height is coded by 3 values: 1 (short), 2 (average), 3 (tall) and shoe size is coded by 3 values 1 (small), 2 (average), 3 (large). The joint counts are given in the following table.

Shoe \ Height	1	2	3
1	234	225	84
2	180	453	161
3	39	192	157

Let X be the coded shoe size and Y the height of a random person in the sample.

- (a) Find the joint and marginal pmf of X and Y .
- (b) Are X and Y independent?

Problem 58. Let X and Y be two continuous random variables with joint pdf

$$f(x, y) = cx^2y(1 + y) \quad \text{for } 0 \leq x \leq 3 \text{ and } 0 \leq y \leq 3,$$

and $f(x, y) = 0$ otherwise.

- (a) Find the value of c .
- (b) Find the probability $P(1 \leq X \leq 2, 0 \leq Y \leq 1)$.
- (c) Determine the joint cdf, $F(a, b)$, of X and Y for a and b between 0 and 3.
- (d) Find marginal cdf $F_X(a)$ for a between 0 and 3.
- (e) Find the marginal pdf $f_X(x)$ directly from $f(x, y)$ and check that it is the derivative of $F_X(x)$.
- (f) Are X and Y independent?

Problem 59. Let X and Y be two random variables and let r , s , t , and u be real numbers.

- (a) Show that $\text{Cov}(X + s, Y + u) = \text{Cov}(X, Y)$.
 (b) Show that $\text{Cov}(rX, tY) = rt\text{Cov}(X, Y)$.
 (c) Show that $\text{Cov}(rX + s, tY + u) = rt\text{Cov}(X, Y)$.

Problem 60. Derive the formula for the covariance: $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$.

Problem 61. (Arithmetic Puzzle)

The joint and marginal pmf's of X and Y are partly given in the following table.

$X \backslash Y$	1	2	3	
1	1/6	0	...	1/3
2	...	1/4	...	1/3
3	1/4	...
	1/6	1/3	...	1

- (a) Complete the table.
 (b) Are X and Y independent?

Problem 62. (Simple Joint Probability)

Let X and Y each have range $\{1, 2, 3, 4\}$. The following formula gives their joint pmf

$$P(X = i, Y = j) = \frac{i + j}{80}$$

Compute each of the following:

- (a) $P(X = Y)$.
 (b) $P(XY = 6)$.
 (c) $P(1 \leq X \leq 2, 2 < Y \leq 4)$.

Problem 63. Toss a fair coin 3 times. Let X = the number of heads on the first toss, Y the total number of heads on the last two tosses, and F the number of heads on the first two tosses.

- (a) Give the joint probability table for X and Y . Compute $\text{Cov}(X, Y)$.
 (b) Give the joint probability table for X and F . Compute $\text{Cov}(X, F)$.

Problem 64. Covariance and Independence

Let X be a random variable that takes values $-2, -1, 0, 1, 2$; each with probability $1/5$. Let $Y = X^2$.

- (a) Fill out the following table giving the joint frequency function for X and Y . Be sure to include the marginal probabilities.

X	-2	-1	0	1	2	total
Y						
0						
1						
4						
total						

- (b) Find $E[X]$ and $E[Y]$.
- (c) Show X and Y are not independent.
- (d) Show $\text{Cov}(X, Y) = 0$.

This is an example of uncorrelated but non-independent random variables. The reason this can happen is that correlation only measures the linear dependence between the two variables. In this case, X and Y are not at all linearly related.

Problem 65. Continuous Joint Distributions

Suppose X and Y are continuous random variables with joint density function $f(x, y) = x + y$ on the unit square $[0, 1] \times [0, 1]$.

- (a) Let $F(x, y)$ be the joint CDF. Compute $F(1, 1)$. Compute $F(x, y)$.
- (b) Compute the marginal densities for X and Y .
- (c) Are X and Y independent?
- (d) Compute $E[X]$, $E[Y]$, $E[X^2 + Y^2]$, $\text{Cov}(X, Y)$.

Problem 66. Correlation

Flip a coin 3 times. Use a joint pmf table to compute the covariance and correlation between the number of heads on the first 2 and the number of heads on the last 2 flips.

Problem 67. Correlation

Flip a coin 5 times. Use properties of covariance to compute the covariance and correlation between the number of heads on the first 3 and last 3 flips.

8 Law of Large Numbers, Central Limit Theorem

Problem 68. (Table of normal probabilities)

Use the table of standard normal probabilities to compute the following. (Z is the standard normal.)

- (a) (i) $P(Z \leq 1.5)$ (ii) $P(-1.5 < Z < 1.5)$ $P(Z > -0.75)$.
- (b) Suppose $X \sim N(2, (0.5)^2)$. Find (i) $P(X \leq 2)$ (ii) $P(1 < X \leq 1.75)$.

Problem 69. Suppose X_1, \dots, X_{100} are i.i.d. with mean $1/5$ and variance $1/9$. Use the central limit theorem to estimate $P(\sum X_i < 30)$.

Problem 70. All or None

You have \$100 and, never mind why, you must convert it to \$1000. Anything less is no good. Your only way to make money is to gamble for it. Your chance of winning one bet is p .

Here are two extreme strategies:

Maximum strategy: bet as much as you can each time. To be smart, if you have less than \$500 you bet it all. If you have more, you bet enough to get to \$1000.

Minimum strategy: bet \$1 each time.

If $p < 0.5$ (the odds are against you) which is the better strategy?

What about $p > 0.5$?

Problem 71. (Central Limit Theorem)

Let X_1, X_2, \dots, X_{81} be i.i.d., each with expected value $\mu = E[X_i] = 5$, and variance $\sigma^2 = \text{Var}(X_i) = 4$. Approximate $P(X_1 + X_2 + \dots + X_{81} > 369)$, using the central limit theorem.

Problem 72. (Binomial \approx normal)

Let $X \sim \text{binomial}(100, 1/3)$.

An 'exact' computation in R gives $P(X \leq 30) = 0.2765539$. Use the central limit theorem to give an approximation of $P(X \leq 30)$

Problem 73. (More Central Limit Theorem)

The average IQ in a population is 100 with standard deviation 15 (by definition, IQ is normalized so this is the case). What is the probability that a randomly selected group of 100 people has an average IQ above 115?

Problem 74. Hospitals (binomial, CLT, etc)

- A certain town is served by two hospitals.
- Larger hospital: about 45 babies born each day.
- Smaller hospital about 15 babies born each day.
- For a period of 1 year, each hospital recorded the days on which more than 60% of the babies born were boys.

(a) Which hospital do you think recorded more such days?

- (i) The larger hospital.
- (ii) The smaller hospital.
- (iii) About the same (that is, within 5% of each other).

(b) Let L_i (resp., S_i) be the Bernoulli random variable which takes the value 1 if more than 60% of the babies born in the larger (resp., smaller) hospital on the i^{th} day were boys. Determine the distribution of L_i and of S_i .

(c) Let L (resp., S) be the number of days on which more than 60% of the babies born in the larger (resp., smaller) hospital were boys. What type of distribution do L and S have? Compute the expected value and variance in each case.

(d) Via the CLT, approximate the 0.84 quantile of L (resp., S). Would you like to revise your answer to part (a)?

(e) What is the correlation of L and S ? What is the joint pmf of L and S ? Visualize the region corresponding to the event $L > S$. Express $P(L > S)$ as a double sum.

9 R Problems

R will not be on the exam. However, these problems will help you understand the concepts we've been studying.

Problem 75. R simulation

Consider X_1, X_2, \dots all independent and with distribution $N(0, 1)$. Let \bar{X}_m be the average of X_1, \dots, X_m .

(a) Give $E[\bar{X}_n]$ and $\sigma_{\bar{X}_n}$ exactly.

(b) Use a R simulation to estimate $E[\bar{X}_n]$ and $\text{Var}(\bar{X}_n)$ for $n = 1, 9, 100$. (You should use the `rnorm` function to simulate 1000 samples of each X_j .)

Problem 76. R Exercise

Let X_1, X_2, X_3, X_4, X_5 be independent $U(0, 1)$ random variables.

Let $X = X_1 + X_2 + X_3$ and $Y = X_3 + X_4 + X_5$.

Use the `runif()` function to simulate 1000 trials of each of these variables. Use these to estimate $\text{Cov}(X, Y)$.

Extra Credit

Compute this covariance exactly.

MIT OpenCourseWare

<https://ocw.mit.edu>

18.05 Introduction to Probability and Statistics

Spring 2022

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.