

CHAPTER 14 MULTIPLE INTEGRALS

14.1 Double Integrals (page 526)

The double integral $\iint_R f(x, y) dA$ gives the volume between R and the surface $z = f(x, y)$. The base is first cut into small squares of area ΔA . The volume above the i th piece is approximately $f(x_i, y_i) \Delta A$. The limit of the sum $\sum f(x_i, y_i) \Delta A$ is the volume integral. Three properties of double integrals are $\iint (f + g) dA = \iint f dA + \iint g dA$ and $\iint c f dA = c \iint f dA$ and $\iint_R f dA = \iint_S f dA + \iint_T f dA$ if R splits into S and T .

If R is the rectangle $0 \leq x \leq 4, 4 \leq y \leq 6$, the integral $\iint x dA$ can be computed two ways. One is $\iint x dy dx$, when the inner integral is $xy|_4^6 = 2x$. The outer integral gives $x^2|_0^4 = 16$. When the x integral comes first it equals $\int x dx = \frac{1}{2}x^2|_0^4 = 8$. Then the y integral equals $8y|_4^6 = 16$. This is the volume between the base rectangle and the plane $z = x$.

The area R is $\iint 1 dy dx$. When R is the triangle between $x = 0, y = 2x$, and $y = 1$, the inner limits on y are $2x$ and 1 . This is the length of a thin vertical strip. The (outer) limits on x are 0 and $\frac{1}{2}$. The area is $\frac{1}{4}$. In the opposite order, the (inner) limits on x are 0 and $\frac{1}{2}y$. Now the strip is horizontal and the outer integral is $\int_0^1 \frac{1}{2}y dy = \frac{1}{4}$. When the density is $\rho(x, y)$, the total mass in the region R is $\iint \rho dx dy$. The moments are $M_y = \iint \rho x dx dy$ and $M_x = \iint \rho y dx dy$. The centroid has $\bar{x} = M_y/M$.

- 1 $\frac{8}{3}, \frac{2}{3}$ 3 $1; \ln \frac{3}{2}$ 5 2 7 $\frac{1}{2}$ 9 $\frac{4}{3}$ 11 $\int_{y=1}^2 \int_{x=1}^2 dx dy + \int_{y=2}^4 \int_{x=y/2}^2 dx dy$
 13 $\int_{y=0}^1 \int_{x=-\frac{1}{2} \ln y}^{-\frac{1}{2} \ln y} dx dy$ 15 $\int_{x=0}^1 \int_{y=-\sqrt{x}}^{\sqrt{x}} dy dx$ 17 $\int_0^1 \int_0^{y/2} dx dy = \int_0^{1/2} \int_{2x}^1 dy dx = \frac{1}{4}$
 19 $\int_0^3 \int_{-y}^y dx dy = \int_{-1}^0 \int_{-x}^3 dy dx + \int_0^1 \int_x^3 dy dx = 9$ 21 $\int_0^4 \int_{y/2}^y dx dy + \int_4^8 \int_{y/2}^4 dx dy = \int_0^4 \int_x^{2x} dy dx = 8$
 23 $\int_0^1 \int_0^{bx} dy dx + \int_1^2 \int_0^{b(2-x)} dy dx = \int_0^b \int_{y/b}^{2-(y/b)} dx dy = b$ 25 $f(a, b) - f(a, 0) - f(0, b) + f(0, 0)$
 27 $\int_0^1 \int_0^1 (2x - 3y + 1) dx dy = \frac{1}{2}$ 29 $\int_a^b f(x) dx = \int_a^b \int_0^{f(x)} 1 dy dx$ 31 50,000π
 33 $\int_1^3 \int_1^2 x^2 dx dy = \frac{14}{3}$ 35 $2 \int_0^1 \int_0^{\sqrt{1-y^2}} 1 dx dy = \frac{\pi}{4}$
 37 $\frac{1}{n^2} \sum_{j=1}^n \sum_{i=1}^n f(\frac{i-1}{n}, \frac{j-1}{n})$ is exact for $f = 1, x, y, xy$ 39 Volume 8.5 41 Volumes $\ln 2, 2 \ln(1 + \sqrt{2})$
 43 $\int_0^1 \int_0^1 x^y dx dy = \int_0^1 \frac{1}{y+1} dy = \ln 2; \int_0^1 \int_0^1 x^y dy dx = \int_0^1 \frac{x-1}{\ln x} dx = \ln 2$
 45 With long rectangles $\sum y_i \Delta A = \sum \Delta A = 1$ but $\iint y dA = \frac{1}{2}$

- 2 $\int_1^e 2xy dx = x^2 y|_1^e = (e^2 - 1)y; \int_2^{2e} (e^2 - 1)y dy = (e^2 - 1) \frac{y^2}{2}|_2^{2e} = (e^2 - 1)(2e^2 - 2) = 2(e^2 - 1)^2;$
 $\int_1^e \frac{dx}{xy} = \frac{\ln x}{y}|_1^e = \frac{1}{y}; \int_2^{2e} \frac{dy}{y} = \ln 2e - \ln 2 = \ln \frac{2e}{2} = 1.$
 4 $\int_1^2 y e^{xy} dx = e^{xy}|_1^2 = e^{2y} - e^y; \int_0^1 (e^{2y} - e^y) dy = [\frac{1}{2}e^{2y} - e^y]|_0^1 = \frac{1}{2}e^2 - e + \frac{1}{2}; \int_0^3 \frac{dy}{\sqrt{3+2x+y}} = 2\sqrt{3+2x+y}|_0^3 =$
 $2\sqrt{6+2x} - 2\sqrt{3+2x};$ the x integral is $[\frac{2}{3}(6+2x)^{3/2} - \frac{2}{3}(3+2x)^{3/2}]_{-1}^1 = \frac{2}{3}8^{3/2} - \frac{2}{3}5^{3/2} - \frac{2}{3}4^{3/2} + \frac{2}{3}.$

Note! $3 + 2x + y$ is not zero in the region of integration.

- 6 The region is above $y = x^3$ and below $y = x$ (from 0 to 1). Area = $\int_0^1 (x - x^3) dx = [\frac{x^2}{2} - \frac{x^4}{4}]_0^1 = \frac{1}{4}.$

- 8 The region is below the parabola $y = 1 - x^2$ and above its mirror image $y = x^2 - 1$.

$$\text{Area} = \int_{-1}^1 (1 - x^2 - x^2 + 1) dx = [2x - \frac{2}{3}x^3]_{-1}^1 = \frac{8}{3}.$$

- 10 The area is all below the axis $y = 0$, where horizontal strips cross from $x = y$ to $x = |y|$ (which is $-y$). Note that the y integral stops at $y = 0$. Area $= \int_{-1}^0 \int_y^{-y} dx dy = \int_{-1}^0 -2y dy = [-y^2]_{-1}^0 = 1$.
- 12 The strips in Problem 6 from $y = x^3$ up to x are changed to strips from $x = y$ across to $x = y^{1/3}$. The outer integral on y is by chance also from 0 to 1. Area $= \int_0^1 (y^{1/3} - y) dy = [\frac{3}{4}y^{4/3} - \frac{1}{2}y^2]_0^1 = \frac{1}{4}$.
- 14 Between the upper parabola $y = 1 - x^2$ in Problem 8 and the x axis, the strips now cross from the left side $x = -\sqrt{1-y}$ to the right side $x = +\sqrt{1-y}$. This half of the area is $\int_0^1 \int_{-\sqrt{1-y}}^{\sqrt{1-y}} dx dy = \int_0^1 2\sqrt{1-y} dy = -\frac{4}{3}(1-y)^{3/2}|_0^1 = \frac{4}{3}$. The other half has strips from left side to right side of $y = x^2 - 1$ or $x = \pm\sqrt{1+y}$. This area is $\int_{-1}^0 \int_{-\sqrt{1+y}}^{\sqrt{1+y}} dx dy$ (also $\frac{4}{3}$).
- 16 The triangle in Problem 10 had sides $x = y$, $x = -y$, and $y = -1$. Now the strips are vertical. They go from $y = -1$ up to $y = x$ on the left side: area $= \int_{-1}^0 \int_{-1}^x dy dx = \int_{-1}^0 (x+1) dx = \frac{1}{2}(x+1)^2|_{-1}^0 = \frac{1}{2}$. The strips go from -1 up to $y = -x$ on the right side: area $= \int_0^1 \int_{-1}^{-x} dy dx = \int_0^1 (-x+1) dx = \frac{1}{2}$. Check: $\frac{1}{2} + \frac{1}{2} = 1$.
- 18 The triangle has corners at $(0,0)$ and $(-1,0)$ and $(-1,-1)$. Its area is $\int_{-1}^0 \int_0^{-x} dy dx = \int_{-1}^0 \int_{-1}^{-y} dx dy (= \frac{1}{2})$.
- 20 The triangle has corners at $(0,0)$ and $(2,4)$ and $(4,4)$. Horizontal strips go from $x = \frac{y}{2}$ to $x = y$: area $= \int_0^4 \int_{y/2}^y dx dy = 4$. Vertical strips are of two kinds: from $y = x$ up to $y = 2x$ or to $y = 4$. Area $= \int_0^2 \int_x^{2x} dy dx + \int_2^4 \int_x^4 dy dx = 2 + 2 = 4$.
- 22 (Hard Problem) The boundary lines are $y = \frac{1}{2}x$ from $(-2,-1)$ to $(0,0)$, and $y = -2x$ from $(0,0)$ to $(1,-2)$, and $y = \frac{-x-5}{3}$ or $x = -3y - 5$ from $(-2,-1)$ to $(1,-2)$. (This is the hardest one: note first the slope $-\frac{1}{3}$.) Vertical strips go from the third line up to the first or second: area $= \int_{-2}^0 \int_{(-x-5)/3}^{x/2} dy dx + \int_0^1 \int_{(-x-5)/3}^{-2x} dy dx = \frac{5}{3} + \frac{5}{6} = \frac{5}{2}$. Horizontal strips cross from the first or third lines to the second: area $= \int_{-2}^{-1} \int_{-3y-5}^{-y/2} dx dy + \int_{-1}^0 \int_{-3y-5}^{-y/2} dx dy = \frac{5}{4} + \frac{5}{4} = \frac{5}{2}$.
- 24 The top of the triangle is (a,b) . From $x = 0$ to a the vertical strips lead to $\int_0^a \int_{dx/c}^{bx/a} dy dx = [\frac{bx^2}{2a} - \frac{dx^2}{2c}]_0^a = \frac{ba}{2} - \frac{da^2}{2c}$. From $x = a$ to c the strips go up to the third side: $\int_a^c \int_{dx/c}^{b+(x-a)(d-b)/(c-a)} dy dx = [bx + \frac{(x-a)^2(d-b)}{2(c-a)} - \frac{dx^2}{2c}]_a^c = b(c-a) + \frac{(c-a)(d-b)}{2} - \frac{dc}{2} + \frac{da^2}{2c}$. The sum is $\frac{ba}{2} + \frac{b(c-a)}{2} + \frac{d(c-a)}{2} - \frac{dc}{2} = \frac{bc-ad}{2}$. This is half of a parallelogram.
- 26 $\int_0^b \int_0^a \frac{\partial f}{\partial x} dx dy = \int_0^b [f(a,y) - f(0,y)] dy$.
- 28 Over the square $\int_0^1 \int_0^1 (xe^y - ye^x) dy dx = \int_0^1 (xe - \frac{e^x}{2} - x) dx = [\frac{x^2 e}{2} - \frac{e^x}{2} - \frac{x^2}{2}]_0^1 = \frac{e}{2} - \frac{e}{2} - \frac{1}{2} + \frac{1}{2} = 0$. (Looking back: zero is not a surprise because of symmetry.) Over the triangle the integral up to $y = x$ is $\int_0^1 \int_0^x (xe^y - ye^x) dy dx$. Over the triangle across to $y = x$ the integral is $\int_0^1 \int_0^y (xe^y - ye^x) dx dy$. Exchange y and x in the second double integral to get minus the first double integral.
- 30 $\int_{-1}^1 (1-x^2) dx = [x - \frac{x^3}{3}]_{-1}^1 = \frac{4}{3}$. With horizontal strips this is $\int_0^1 \int_{-\sqrt{1-y}}^{\sqrt{1-y}} dx dy = \int_0^1 2\sqrt{1-y} dy = -\frac{4}{3}(1-y)^{3/2}|_0^1 = \frac{4}{3}$.
- 32 The height is $z = \frac{1-ax-by}{c}$. Integrate over the triangular base ($z = 0$ gives the side $ax + by = 1$): volume $= \int_{x=0}^{1/a} \int_{y=0}^{(1-ax)/b} \frac{1-ax-by}{c} dy dx = \int_0^{1/a} [y - axy - \frac{1}{2}by^2]_0^{(1-ax)/b} dx = \int_0^{1/a} \frac{1}{c} \frac{(1-ax)^2}{2b} dx = -\frac{(1-ax)^3}{6abc} \Big|_0^{1/a} = \frac{1}{6abc}$.
- 34 From Problem 33 the mass is $\frac{14}{3}$. The moments are $\int_1^3 \int_1^2 x^3 dx dy = \int_1^3 \frac{2^4-1^4}{4} dy = \frac{15}{2}$ and $\int_1^3 \int_1^2 yx^2 dx dy = \int_1^3 \frac{8-1}{3} y dy = \frac{28}{3}$. Then $\bar{x} = \frac{15/2}{14/3} = \frac{45}{28}$ and $\bar{y} = \frac{28/3}{14/3} = 2$.
- 36 The area of the quarter-circle is $\frac{\pi}{4}$. The moment is zero around the axis $y = 0$ (by symmetry): $\bar{x} = 0$. The other moment, with a factor 2 that accounts for symmetry of left and right, is $2 \int_0^{\sqrt{2}/2} \int_x^{\sqrt{1-x^2}} y dy dx = 2 \int_0^1 (\frac{1-x^2}{2} - \frac{x^2}{2}) dx = 2[\frac{x}{2} - \frac{x^3}{3}]_0^{\sqrt{2}/2} = \frac{\sqrt{2}}{3}$. Then $\bar{y} = \frac{\sqrt{2}/3}{\pi/4} = \frac{4\sqrt{2}}{3\pi}$.
- 38 The integral $\int_0^1 \int_0^1 x^2 dx dy$ has the usual midpoint error $-\frac{(\Delta x)^2}{12}$ for the integral of x^2 (see Section 5.8). The y integral $\int_0^1 dy = 1$ is done exactly. So the error is $-\frac{1}{12n^2}$ (and the same for $\iint y^2 dx dy$). The integral of xy is computed exactly. Errors decrease with exponent $p = 2$, the order of accuracy.

40 The exact integral is $\int_0^1 \int_0^1 \frac{dx dy}{\sqrt{x^2+y^2}} = 2 \int_0^{\pi/4} \int_0^{\sec \theta} \frac{r dr d\theta}{r} = 2 \int_0^{\pi/4} \sec \theta d\theta = 2 \ln(\sec \theta + \tan \theta) \Big|_0^{\pi/4} = 2 \ln(\sqrt{2} + 1).$

42 The exact integral is $\int_0^1 \int_0^1 e^x \sin \pi y dx dy = \int_0^1 (e - 1) \sin \pi y dy = \frac{e-1}{\pi} (-\cos \pi y) \Big|_0^1 = \frac{2}{\pi} (e - 1).$

14.2 Change to Better Coordinates (page 534)

We change variables to improve the limits of integration. The disk $x^2 + y^2 \leq 9$ becomes the rectangle $0 \leq r \leq 3, 0 \leq \theta \leq 2\pi$. The inner limits of $\iint dy dx$ are $y = \pm\sqrt{9-x^2}$. In polar coordinates this area integral becomes $\iint r dr d\theta = 9\pi$.

A polar rectangle has sides dr and $r d\theta$. Two sides are not straight but the angles are still 90° . The area between the circles $r = 1$ and $r = 3$ and the rays $\theta = 0$ and $\theta = \pi/4$ is $\frac{1}{8}(3^2 - 1^2) = 1$. The integral $\iint x dy dx$ changes to $\iint r^2 \cos \theta dr d\theta$. This is the moment around the y axis. Then $\bar{x}$ is the ratio M_y/M . This is the x coordinate of the centroid, and it is the average value of x .

In a rotation through α , the point that reaches (u, v) starts at $x = u \cos \alpha - v \sin \alpha, y = u \sin \alpha + v \cos \alpha$. A rectangle in the uv plane comes from a rectangle in xy . The areas are equal so the stretching factor is $J = 1$. This is the determinant of the matrix $\begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$. The moment of inertia $\iint x^2 dx dy$ changes to $\iint (u \cos \alpha - v \sin \alpha)^2 du dv$.

For single integrals dx changes to $(dx/du)du$. For double integrals $dx dy$ changes to $J du dv$ with $J = \partial(x, y)/\partial(u, v)$. The stretching factor J is the determinant of the 2 by 2 matrix $\begin{bmatrix} \partial x/\partial u & \partial x/\partial v \\ \partial y/\partial u & \partial y/\partial v \end{bmatrix}$. The functions $x(u, v)$ and $y(u, v)$ connect an xy region R to a uv region S , and $\iint_R dx dy = \iint_S J du dv = \text{area of } R$. For polar coordinates $x = u \cos v$ and $y = u \sin v$ (or $r \sin \theta$). For $x = u, y = u + 4v$ the 2 by 2 determinant is $J = 4$. A square in the uv plane comes from a parallelogram in xy . In the opposite direction the change has $u = x$ and $v = \frac{1}{4}(y - x)$ and a new $J = \frac{1}{4}$. This J is constant because this change of variables is linear.

1 $\int_{\pi/4}^{3\pi/4} \int_0^1 r dr d\theta = \frac{\pi}{4}$ 3 S = quarter-circle with $u \geq 0$ and $v \geq 0$; $\int_0^1 \int_0^{\sqrt{1-v^2}} du dv$

5 R is symmetric across the y axis; $\int_0^1 \int_0^{\sqrt{1-v^2}} u du dv = \frac{1}{3}$ divided by area gives $(\bar{u}, \bar{v}) = (4/3\pi, 4/3\pi)$

7 $2 \int_0^{1/\sqrt{2}} \int_{1+x}^{1+\sqrt{1-x^2}} dy dx$; xy region R^* becomes R in the x^*y^* plane; $dx dy = dx^* dy^*$ when region moves

9 $J = \begin{vmatrix} \partial x/\partial r^* & \partial x/\partial \theta^* \\ \partial y/\partial r^* & \partial y/\partial \theta^* \end{vmatrix} = \begin{vmatrix} \cos \theta^* & -r^* \sin \theta^* \\ \sin \theta^* & r^* \cos \theta^* \end{vmatrix} = r^*$; $\int_{\pi/4}^{3\pi/4} \int_0^1 r^* dr^* d\theta^*$

11 $I_y = \iint_R x^2 dx dy = \int_{\pi/4}^{3\pi/4} \int_0^1 r^2 \cos^2 \theta r dr d\theta = \frac{\pi}{16} - \frac{1}{8}$; $I_x = \frac{\pi}{16} + \frac{1}{8}$; $I_0 = \frac{\pi}{8}$

13 $(0,0), (1,2), (1,3), (0,1)$; area of parallelogram is 1

15 $x = u, y = u + 3v + uv$; then $(u, v) = (1,0), (1,1), (0,1)$ give corners $(x, y) = (1,0), (1,5), (0,3)$

17 Corners $(0,0), (2,1), (3,3), (1,2)$; sides $y = \frac{1}{2}x, y = 2x - 3, y = \frac{1}{2}x + \frac{3}{2}, y = 2x$

19 Corners $(1,1), (e^2, e), (e^3, e^3), (e, e^2)$; sides $x = y^2, y = x^2/e^3, x = y^2/e^3, y = x^2$

21 Corners $(0,0), (1,0), (1,2), (0,1)$; sides $y = 0, x = 1, y = 1 + x^2, x = 0$

$$23 \quad J = \begin{vmatrix} 2 & 1 \\ 1 & 2 \end{vmatrix} = 3, \text{ area } \int_0^1 \int_0^1 3 \, du \, dv = 3; J = \begin{vmatrix} 2e^{2u+v} & e^{2u+v} \\ e^{u+2v} & 2e^{u+2v} \end{vmatrix} = 3e^{3u+3v}, \int_0^1 \int_0^1 3e^{3u+3v} \, du \, dv = \int_0^1 (e^{3+3v} - e^{3v}) \, dv = \frac{1}{3}(e^6 - e^3 + 1)$$

$$25 \quad \text{Corners } (x, y) = (0, 0), (1, 0), (1, f(1)), (0, f(0)); (\frac{1}{2}, 1) \text{ gives } x = \frac{1}{2}, y = f(\frac{1}{2}); J = \begin{vmatrix} 1 & 0 \\ v f'(u) & f(u) \end{vmatrix} = f(u)$$

$$27 \quad B^2 = 2 \int_0^{\pi/4} \int_0^{1/\sin \theta} e^{-r^2} r \, dr \, d\theta = \int_0^{\pi/4} (e^{-1/\sin^2 \theta} - 1) d\theta$$

$$29 \quad \bar{r} = \iint r^2 \, dr \, d\theta / \iint r \, dr \, d\theta = \int_0^\pi \frac{8}{3} a^3 \sin^3 \theta \, d\theta / \pi a^2 = \frac{32a}{9\pi} \quad 31 \quad \int_0^{2\pi} \int_0^1 r^2 r \, dr \, d\theta = \frac{\pi}{2}$$

33 Along the right side; along the bottom; at the bottom right corner

$$35 \quad \iint xy \, dx \, dy = \int_0^1 \int_0^1 (u \cos \alpha - v \sin \alpha)(u \sin \alpha + v \cos \alpha) \, du \, dv = \frac{1}{4}(\cos^2 \alpha - \sin^2 \alpha)$$

$$37 \quad \int_0^{2\pi} \int_4^5 r^2 r^2 \, dr \, d\theta = \frac{2\pi}{6}(5^6 - 4^6) \quad 39 \quad x = \cos \alpha - \sin \alpha, y = \sin \alpha + \cos \alpha \text{ goes to } u = 1, v = 1$$

$$2 \quad \text{Area} = \int_{-\sqrt{2}/2}^{\sqrt{2}/2} \int_{|x|}^{\sqrt{1-x^2}} dy \, dx \text{ splits into two equal parts left and right of } x = 0: 2 \int_0^{\sqrt{2}/2} \int_x^{\sqrt{1-x^2}} dy \, dx = 2 \int_0^{\sqrt{2}/2} (\sqrt{1-x^2} - x) \, dx = [x\sqrt{1-x^2} + \sin^{-1} x - x^2]_0^{\sqrt{2}/2} = \sin^{-1} \frac{\sqrt{2}}{2} = \frac{\pi}{4}. \text{ The limits on}$$

$$\iint dx \, dy \text{ are } \int_0^{\sqrt{2}/2} \int_{-y}^y dx \, dy \text{ for the lower triangle plus } \int_{\sqrt{2}/2}^1 \int_{-\sqrt{1-y^2}}^{\sqrt{1-y^2}} dx \, dy \text{ for the circular top.}$$

$$4 \quad (\text{See Problem 36 of Section 14.1}) \int_{\pi/4}^{3\pi/4} \int_0^1 (r \sin \theta) r \, dr \, d\theta = [\frac{r^2}{2}]_0^1 [-\cos \theta]_{\pi/4}^{3\pi/4} = \frac{\sqrt{2}}{3}; \text{ divide by area } \frac{\pi}{4} \text{ to reach } \bar{y} = \frac{\sqrt{2}/3}{\pi/4} = \frac{4\sqrt{2}}{3\pi}.$$

$$6 \quad \text{Area of wedge} = \frac{b}{2\pi}(\pi a^2). \text{ Divide } \int_0^b \int_0^a (r \cos \theta) r \, dr \, d\theta = \frac{a^3}{3} \sin b \text{ by this area } \frac{ba^2}{2} \text{ to find } \bar{x} = \frac{2a}{3b} \sin b. (\text{Interesting limit: } \bar{x} \rightarrow \frac{2}{3}a \text{ as the wedge angle } b \text{ approaches zero: like the centroid of a triangle.})$$

$$\text{For } \bar{y} \text{ divide } \int_0^b \int_0^a (r \sin \theta) r \, dr \, d\theta = \frac{a^3}{3} (1 - \cos b) \text{ by the area } \frac{ba^2}{2} \text{ to find } \bar{y} = \frac{2a}{3b} (1 - \cos b).$$

8 The limits on r, θ are extremely awkward for R^* . Contrast with the simple limits $0 \leq r^* \leq 1, \frac{\pi}{4} \leq \theta^* \leq \frac{3\pi}{4}$ when the coordinates are recentered at $(0, 1)$. (A point on the lower boundary of the wedge has $r = \frac{\sin \frac{3\pi}{4}}{\sin(\frac{\pi}{4} - \theta)}$ by the law of sines.)

10 The centroid $(0, \bar{y})$ of R moves up to the centroid $(0, \bar{y} + 1)$ of R^* . The centroid of a circle is its center $(1, 2)$. The centroid of the upper half is $(1, 2 + \frac{4}{\pi})$ because a half-circle has $\int_0^\pi \int_0^3 (r \sin \theta) r \, dr \, d\theta = 18$ divided by its area $\frac{9\pi}{2}$ (which gives $\frac{4}{\pi}$).

$$12 \quad I_x = \int_{\pi/4}^{3\pi/4} \int_0^1 (r \sin \theta + 1)^2 r \, dr \, d\theta = \frac{1}{4} \int \sin^2 \theta \, d\theta + \frac{2}{3} \int \sin \theta \, d\theta + \frac{1}{2} \int d\theta = [\frac{\theta}{8} - \frac{\sin 2\theta}{16} - \frac{2}{3} \cos \theta + \frac{\theta}{2}]_{\pi/4}^{3\pi/4} = \frac{5\pi}{16} + \frac{2}{16} + \frac{4}{3} \frac{\sqrt{2}}{2}; I_y = \iint (r \cos \theta)^2 r \, dr \, d\theta = \frac{\pi}{16} - \frac{1}{8} \text{ (as in Problem 11); } I_0 = I_x + I_y = \frac{3\pi}{8} + \frac{4}{3} \frac{\sqrt{2}}{2}.$$

14 The corner $(1, 2)$ should be (a, c) , when $u = 0$ and $v = 1$; the corner $(0, 1)$ should be (b, d) , when $u = 1$ and $v = 0$. Check at $u = v = 1$; there $x = au + bv = 1$ and $y = cu + dv = 3$ to give the correct corner $(1, 3)$.

Then $J = ad - bc = (1)(1) - (0)(2) = 1$. The unit square has area 1; so does R .

16 A linear change takes the square S into a parallelogram R (with one corner at $(0, 0)$). Reason: The vector sum of the two sides from $(0, 0)$ is still the vector to the far corner.

18 Corners when $u = 0$ or $1, v = 0$ or 1 : $(0, 0), (3, 1), (5, 2), (2, 1)$. The sides have equations

$$y = \frac{1}{3}x, y = \frac{1}{2}x - \frac{1}{2}, y = \frac{1}{3}x + \frac{1}{3}, y = \frac{1}{2}x.$$

20 Corners when $u = 0$ or $1, v = 0$ or 1 : $(0, 0), (0, -1), (1, 0), (0, 1)$. Actually $(0, 0)$ is not a corner because one side comes down the y axis. The side with $u = 1$ is $x = v, y = v^2 - 1$ or $y = x^2 - 1$. The side with $v = 1$ is $x = u, y = 1 - u^2$ or $y = 1 - x^2$.

22 Here $u = 0$ or $1, v = 0$ or 1 gives the corners $(0, 0), (1, 0), (\cos 1, \sin 1)$. The side with $u = 1$ is a circular arc $x = \cos v, y = \sin v$ between the last two corners. The other sides are straight: the region is pie-shaped (a fraction $\frac{1}{2\pi}$ of the unit circle).

24 Problem 18 has $J = \begin{vmatrix} 3 & 2 \\ 1 & 1 \end{vmatrix} = 1$. So the area of R is $1 \times$ area of unit square $= 1$. Problem 20 has

$J = \begin{vmatrix} v & u \\ -2u & 2v \end{vmatrix} = 2(u^2 + v^2)$, and integration over the square gives area of $R =$

$\int_0^1 \int_0^1 2(u^2 + v^2) du dv = \frac{4}{3}$. Check in x, y coordinates: area of $R = 2 \int_0^1 (1 - x^2) dx = \frac{4}{3}$.

26 $\begin{vmatrix} \partial r / \partial x & \partial r / \partial y \\ \partial \theta / \partial x & \partial \theta / \partial y \end{vmatrix} = \begin{vmatrix} x/r & y/r \\ -y/r^2 & x/r^2 \end{vmatrix} = \frac{x^2 + y^2}{r^3} = \frac{1}{r}$. As in equation 12, this new J is $\frac{1}{\text{old } J}$.

28 $\int_{-\infty}^{\infty} x^2 e^{-x^2/2} dx = (u)(v) - \int v du = (x)(-e^{-x^2/2}) \Big|_{-\infty}^{\infty} + \int_{-\infty}^{\infty} e^{-x^2/2} dx = 0 + \sqrt{2\pi}$ by Example 5. Divide by $\sqrt{2\pi}$ to find $\sigma^2 = 1$.

30 R is an infinite strip above the interval $[0, 1]$ on the x axis. Its boundary $x = 1$ is $r \cos \theta = 1$ or $r = \sec \theta$. The limits are $0 \leq r \leq \sec \theta$ and $0 \leq \theta \leq \frac{\pi}{2}$. The integral is $\int_0^{\pi/2} \int_0^{\sec \theta} \frac{r}{r^3} dr d\theta = \int_0^{\pi/2} (\infty) d\theta = \text{infinite}$.

For a finite example integrate $(x^2 + y^2)^{-1/2} = \frac{1}{r}$.

32 Equation (3) with y instead of x has $\iint y^2 dA = \int_0^1 \int_0^1 (u \sin \alpha + v \cos \alpha)^2 du dv = \sin^2 \alpha \iint u^2 du dv + \sin \alpha \cos \alpha \iint 2uv du dv + \cos^2 \alpha \iint v^2 du dv = \frac{\sin^2 \alpha}{3} + \frac{\sin \alpha \cos \alpha}{2} + \frac{\cos^2 \alpha}{3}$.

34 (a) **False** (forgot the stretching factor J) (b) **False** (x can be larger than x^2) (c) **False** (forgot to divide by the area) (d) **True** (odd function integrated over symmetric interval) (e) **False** (the straight-sided region is a trapezoid: angle from 0 to θ and radius from r_1 to r_2 yields area $\frac{1}{2}(r_2^2 - r_1^2) \sin \theta \cos \theta$).

36 $\iint \rho dA = \int_0^{2\pi} \int_4^5 r^2 (r dr d\theta) = 2\pi \frac{5^4 - 4^4}{4}$. This is the polar moment of inertia I_0 with density $\rho = 1$.

38 $\iint f dA = f(P) \iint dA$ is the Mean Value Theorem for double integrals (compare Property 7, Section 5.6). If $f = x$ or $f = y$, choose $P = \text{centroid } (\bar{x}, \bar{y})$.

14.3 Triple Integrals (page 540)

Six important solid shapes are a **box**, **prism**, **cone**, **cylinder**, **tetrahedron**, and **sphere**. The integral $\iiint dx dy dz$ adds the volume $dx dy dz$ of small boxes. For computation it becomes three single integrals. The inner integral $\int dx$ is the **length** of a line through the solid. The variables y and z are held constant. The double integral $\iint dx dy$ is the **area** of a slice, with z held constant. Then the z integral adds up the volumes of **slices**.

If the solid region V is bounded by the planes $x = 0, y = 0, z = 0$, and $x + 2y + 3z = 1$, the limits on the inner x integral are **0** and **$1 - 2y - 3z$** . The limits on y are **0** and **$\frac{1}{2}(1 - 3z)$** . The limits on z are **0** and **$\frac{1}{3}$** . In the new variables $u = x, v = 2y, w = 3z$, the equation of the outer boundary is $u + v + w = 1$. The volume of the tetrahedron in uvw space is $\frac{1}{6}$. From $dx = du$ and $dy = dv/2$ and $dz = dw/3$, the volume of an xyz box is $dx dy dz = \frac{1}{6} du dv dw$. So the volume of V is $\frac{1}{36}$.

To find the average height $\bar{z}$ in V we compute $\iiint z dV / \iiint dV$. To find the total mass if the density is $\rho = e^z$ we compute the integral $\iiint e^z dx dy dz$. To find the average density we compute $\iiint e^z dV / \iiint dV$. In the order $\iiint dz dx dy$ the limits on the inner integral can depend on x and y . The limits on the middle integral can depend on y . The outer limits for the ellipsoid $x^2 + 2y^2 + 3z^2 \leq 8$ are $-2 \leq y \leq 2$.

$$1 \int_0^1 \int_0^z \int_0^y dx dy dz = \frac{1}{6}$$

3 $0 \leq y \leq x \leq z \leq 1$ and all other orders xyz, yzx, zxy, zyx ; all six contain $(0, 0, 0)$; to contain $(1, 0, 1)$

- 5 $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 8$ 7 $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 4$ 9 $\int_{-1}^1 \int_x^1 \int_1^x dx dy dz = \frac{4}{3}$
 11 $\int_0^1 \int_0^{2-2z} \int_0^{2-y-2z} dx dy dz = \frac{2}{3}$ 13 $\int_0^{1/3} \int_0^{2-2z} \int_0^{2-y-2z} dx dy dz = \frac{7}{12}$
 15 $\int_0^1 \int_0^{1-z} \int_0^{\sqrt{(1-z)^2-y^2}} dx dy dz = \frac{\pi}{3}$ 17 $\int_0^6 \int_0^1 \int_0^{\sqrt{1-y^2}} dx dy dz = 6\pi$ 19 $\int_0^1 \int_0^1 \int_0^{\sqrt{1-y^2}} dx dy dz = \pi$
 21 Corner of cube at $(\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$; sides $\frac{2}{\sqrt{3}}$; area $\frac{8}{3\sqrt{3}}$
 23 Horizontal slices are circles of area $\pi r^2 = \pi(4-z)$; volume $= \int_0^4 \pi(4-z) dz = 8\pi$; centroid
 has $\bar{x} = 0, \bar{y} = 0, \bar{z} = \int_0^4 z\pi(4-z) dz / 8\pi = \frac{4}{3}$
 25 $I = \frac{z^2}{2}$ gives zeros; $\frac{\partial I}{\partial x} = \int_0^x \int_0^y f dy dz, \frac{\partial I}{\partial y} = \int_0^x \int_0^x f dx dz, \frac{\partial^2 I}{\partial y \partial z} = \int_0^x f dx$
 27 $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 (y^2 + z^2) dx dy dz = \frac{16}{3}$; $\iiint x^2 dV = \frac{8}{3}$; $3 \iiint (x - \frac{x+y+z}{3})^2 dV = \frac{16}{3}$
 29 $\int_0^3 \int_0^2 \int_0^y dx dy dz = 6$ 31 Trapezoidal rule is second-order; correct for 1, x, y, z, xy, xz, yz, xyz

- 2 The area of $0 \leq x \leq y \leq z \leq 1$ is $\int_0^1 \int_x^1 \int_y^1 dz dy dx$. The four faces are $x=0, y=x, z=y, z=1$.
 4 $\int_0^1 \int_0^x \int_0^y x dx dy dz = \int_0^1 \int_0^x \frac{y^2}{2} dy dz = \int_0^1 \frac{z^3}{6} dz = \frac{1}{24}$. Divide by the volume $\frac{1}{6}$ to find $\bar{x} = \frac{1}{4}$;
 $\int_0^1 \int_0^x \int_0^y y dx dy dz = \int_0^1 \int_0^x \frac{y^2}{2} dy dz = \int_0^1 \frac{z^3}{6} dz = \frac{1}{24}$ and $\bar{y} = \frac{1}{2}$; by symmetry $\bar{z} = \frac{3}{4}$.
 6 Volume of half-cube $= \int_0^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = 4$.
 8 $\int_0^1 \int_{-1}^1 \int_{-1}^1 dz dy dx = \int_0^1 2(z+1) dz = [(z+1)^2]_0^1 = 3$.
 10 $\int_{-1}^1 \int_{-1}^1 \int_{-1}^1 dx dy dz = \int_{-1}^1 \int_{-1}^1 (y+1) dy dz = \int_{-1}^1 \frac{(z+1)^2}{2} dz = [\frac{(z+1)^3}{6}]_{-1}^1 = \frac{4}{3}$ (tetrahedron).
 12 The plane faces are $x=0, y=0, z=0$, and $2x+y+z=4$ (which goes through 3 points). The volume
 is $\int_0^2 \int_0^{4-2x} \int_0^{4-2x-y} dz dy dx = \int_0^2 \int_0^{4-2x} (4-2x-y) dy dx = \int_0^2 \frac{(4-2x)^2}{2} dx = [-\frac{(4-2x)^3}{12}]_0^2 = \frac{4^3}{12} = \frac{16}{3}$.
 Check: Multiply standard volume $\frac{1}{6}$ by $(4)(4)(2) = \frac{16}{3}$. Check: Double the volume in Problem 11.
 14 Put dz last and stop at $z=1$: $\int_0^1 \int_0^{4-z} \int_0^{(4-y-z)/2} dx dy dz = \int_0^1 \int_0^{4-z} \frac{4-y-z}{2} dy dz =$
 $\int_0^1 \frac{(4-z)^2}{4} dz = [-\frac{(4-z)^3}{12}]_0^1 = \frac{4^3-3^3}{12} = \frac{37}{12}$.
 16 (Still tetrahedron of Problem 12: volume still $\frac{16}{3}$). Limits of integration: the top vertex
 falls from $(0,0,4)$ onto the y axis at $(0,-4,0)$. The corner $(2,0,0)$ stays on the x axis.
 The corner $(0,4,0)$ swings up to $(0,0,4)$. The volume integral is $\int_0^4 \int_{-4}^0 \int_0^2 dx dy dz = \frac{16}{3}$.
 18 The plane $z=x$ cuts the circular base in half, leaving $x \geq 0$. Volume $= \int_0^1 \int_{-\sqrt{1-x^2}}^{\sqrt{1-x^2}} \int_0^x dz dy dx =$
 $\int_0^1 2x\sqrt{1-x^2} dx = [-\frac{2}{3}(1-x^2)^{3/2}]_0^1 = \frac{2}{3}$.
 20 Lying along the x axis the cylinder goes from $x=0$ to $x=6$. Its slices are circular disks $y^2 + (z-1)^2 = 1$
 resting on the x axis. Volume $= \int_0^6 \int_{-1-\sqrt{1-y^2}}^{1+\sqrt{1-y^2}} dz dy dx =$ still 6π .
 22 Change variables to $X = \frac{x}{a}, Y = \frac{y}{b}, Z = \frac{z}{c}$; then $dXdYdZ = \frac{dxdydz}{abc}$. Volume $= \iiint abc dXdYdZ =$
 $\frac{1}{6}abc$. Centroid $(\bar{x}, \bar{y}, \bar{z}) = (a\bar{X}, b\bar{Y}, c\bar{Z}) = (\frac{a}{4}, \frac{b}{4}, \frac{c}{4})$. (Recall volume $\frac{1}{6}$ and centroid $(\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ of standard
 tetrahedron: this is Example 2.)
 24 (a) Change variables to $X = \frac{x}{4}, Y = \frac{y}{2}, Z = \frac{3z}{4}$. Then the solid is $X^2 + Y^2 + Z^2 = 1$, a unit sphere of volume
 $\frac{4\pi}{3}$. Therefore the original volume is $\frac{4\pi}{3}(4)(2)(\frac{4}{3}) = \frac{128\pi}{9}$. (b) The hypervolume in 4 dimensions is $\frac{1}{24}$,
 following the pattern of 1 for interval, $\frac{1}{2}$ for triangle, $\frac{1}{6}$ for tetrahedron.
 26 Average of $f = \iiint f(x,y,z) dV / \iiint dV =$ integral of $f(x,y,z)$ divided by the volume.
 28 Volume of unit cube $= \sum_{i=1}^{1/\Delta x} \sum_{j=1}^{1/\Delta x} \sum_{k=1}^{1/\Delta x} (\Delta x)^3 = 1$.
 30 In one variable, the midpoint rule is correct for the functions 1 and x . In three variables it is correct for
 1, x, y, z, xy, xz, yz, xyz .
 32 Simpson's Rule has coefficients $\frac{1}{6}, \frac{4}{6}, \frac{1}{6}$ over a unit interval. In three dimensions the 8 corners of the cube will
 have coefficients $(\frac{1}{6})^3 = \frac{1}{216}$. The center will have $(\frac{4}{6})^3 = \frac{64}{216}$. The centers of the 12 edges will have
 $(\frac{1}{6})^2(\frac{4}{6}) = \frac{4}{216}$. The centers of the 6 faces have $(\frac{1}{6})(\frac{4}{6})^2 = \frac{16}{216}$. (Check: $8(1) + 64 + 12(4) + 6(16) = 216$).
 When N^3 cubes are stacked together, with N small cubes each way, there are only $2N + 1$ meshpoints

along each direction. This makes $(2N + 1)^3$ points or about 8 per cube. (Visualize the 8 new points of the cube as having x, y, z equal to zero or $\frac{1}{2}$.)

14.4 Cylindrical and Spherical Coordinates (page 547)

The three cylindrical coordinates are $r\theta z$. The point at $x = y = z = 1$ has $r = \sqrt{2}, \theta = \pi/4, z = 1$. The volume integral is $\iiint r \, dr \, d\theta \, dz$. The solid region $1 \leq r \leq 2, 0 \leq \theta \leq 2\pi, 0 \leq z \leq 4$ is a hollow cylinder (a pipe). Its volume is 12π . From the r and θ integrals the area of a ring (or washer) equals 3π . From the z and θ integrals the area of a shell equals $2\pi rz$. In $r\theta z$ coordinates the shapes of cylinders are convenient, while boxes are not.

The three spherical coordinates are $\rho\phi\theta$. The point at $x = y = z = 1$ has $\rho = \sqrt{3}, \phi = \cos^{-1} 1/\sqrt{3}, \theta = \pi/4$. The angle ϕ is measured from the z axis. θ is measured from the x axis. ρ is the distance to the origin, where r was the distance to the z axis. If $\rho\phi\theta$ are known then $x = \rho \sin \phi \cos \theta, y = \rho \sin \phi \sin \theta, z = \rho \cos \phi$. The stretching factor J is a 3 by 3 determinant and volume is $\iiint r^2 \sin \phi \, dr \, d\phi \, d\theta$.

The solid region $1 \leq \rho \leq 2, 0 \leq \phi \leq \pi, 0 \leq \theta \leq 2\pi$ is a hollow sphere. Its volume is $4\pi(2^3 - 1^3)/3$. From the ϕ and θ integrals the area of a spherical shell at radius ρ equals $4\pi\rho^2$. Newton discovered that the outside gravitational attraction of a sphere is the same as for an equal mass located at the center.

- 1 $(r, \theta, z) = (D, 0, 0); (\rho, \phi, \theta) = (D, \frac{\pi}{2}, 0)$ 3 $(r, \theta, z) = (0, \text{any angle}, D); (\rho, \phi, \theta) = (D, 0, \text{any angle})$
 5 $(x, y, z) = (2, -2, 2\sqrt{2}); (r, \theta, z) = (2\sqrt{2}, -\frac{\pi}{4}, 2\sqrt{2})$ 7 $(x, y, z) = (0, 0, -1); (r, \theta, z) = (0, \text{any angle}, -1)$
 9 $\phi = \tan^{-1}(\frac{z}{r})$ 11 45° cone in unit sphere: $\frac{2\pi}{3}(1 - \frac{1}{\sqrt{2}})$ 13 cone without top: $\frac{7\pi}{3}$
 15 $\frac{1}{4}$ hemisphere: $\frac{\pi}{6}$ 17 $\frac{\pi^2}{8}$ 19 Hemisphere of radius π : $\frac{2}{3}\pi^4$ 21 $\pi(R^2 - z^2); 4\pi r\sqrt{R^2 - r^2}$
 23 $\frac{2}{3}a^3 \tan \alpha$ (see 8.1.39) 27 $\frac{\partial q}{\partial D} = \frac{\rho - D \cos \phi}{q} = \frac{\text{near side}}{\text{hypotenuse}} = \cos \alpha$
 31 Wedges are not exactly similar; the error is higher order $\Rightarrow$ proof is correct
 33 Proportional to $1 + \frac{1}{h}(\sqrt{a^2 + (D-h)^2} - \sqrt{a^2 + D^2})$
 35 $J = \begin{vmatrix} a & & \\ & b & \\ & & c \end{vmatrix} = abc$; straight edges at right angles 37 $\begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r$
 39 $\frac{8\pi\rho^4}{3}; \frac{2}{3}$ 41 $\rho^3; \rho^2$; force = 0 inside hollow sphere

- 2 $(r, \theta, z) = (D, \frac{3\pi}{2}, 0); (\rho, \phi, \theta) = (D, \frac{\pi}{2}, \frac{3\pi}{2})$ 4 $(r, \theta, z) = (5, \cos^{-1} \frac{3}{5}, 5); (\rho, \phi, \theta) = (5\sqrt{2}, \frac{\pi}{4}, \cos^{-1} \frac{3}{5})$
 6 $(x, y, z) = (\frac{3}{2}, \frac{\sqrt{3}}{2}, 1); (r, \theta, z) = (\sqrt{3}, \frac{\pi}{6}, 1)$ 8 $x = r$ on the positive x axis ($x \geq 0, y = 0 (= \theta), z = 0$)
 10 $x = \cos t, y = \frac{\sqrt{2}}{2} \sin t, z = \frac{\sqrt{2}}{2} \sin t$. The unit sphere intersects the plane $y = z$.
 12 The surface $z = 1 + r^2 = 1 + x^2 + y^2$ is a paraboloid (parabola rotated around the z axis). The region is above the half-disk $0 \leq r \leq 1, 0 \leq \theta \leq \pi$. The volume is $\frac{3}{4}\pi$.
 14 This is the volume of a half-cylinder (because of $0 \leq \theta \leq \pi$): height π , radius π , volume $\frac{1}{2}\pi^4$.
 16 The upper surface $\rho = 2$ is the top of a sphere. The lower surface $\rho = \sec \phi$ is the plane $z = \rho \cos \phi = 1$. (The angle $\phi = \frac{\pi}{3}$ is the meeting of sphere and plane, where $\sec \phi = 2$.) The volume is $2\pi \int_0^{\pi/3} (\frac{8 - \sec^3 \phi}{3}) \sin \phi \, d\phi = 2\pi [-\frac{8}{3} \cos \phi - \frac{1}{6 \cos^2 \phi}]_0^{\pi/3} = 2\pi [-\frac{4}{3} - \frac{1}{6/4} + \frac{8}{3} + \frac{1}{6}] = \frac{5\pi}{3}$.

- 18** The region $1 \leq \rho \leq 3$ is a hollow sphere (spherical shell). The limits $0 \leq \phi \leq \frac{\pi}{4}$ keep the part that lies above a 45° cone. The volume is $\frac{52\pi}{3}(1 - \frac{\sqrt{2}}{2})$.
- 20** From the unit ball $\rho \leq 1$ keep the part above the cone $\phi = 1$ radian and inside the wedge $0 \leq \theta \leq 1$ radian. Volume $= \frac{1}{4} \int_0^1 \sin \phi d\phi = \frac{1}{4}(1 - \cos 1)$.
- 22** The curve $\rho = 1 - \cos \phi$ is a cardioid in the xz plane (like $r = 1 - \cos \theta$ in the xy plane). So we have a **cardioid of revolution**. Its volume is $\frac{8\pi}{3}$ as in Problem 9.3.35.
- 24** Mass $= \int_0^{2\pi} \int_0^\pi \int_0^R \rho \sin \phi (\rho + 1) d\rho d\phi d\theta = \frac{4}{3}\pi R^3 + 2\pi R^2$.
- 26** *Newton's achievement* The cosine law (see hint) gives $\cos \alpha = \frac{D^2 + q^2 - \rho^2}{2qD}$. Then integrate $\frac{\cos \alpha}{q^3}$: $\iiint (\frac{D^2 - \rho^2}{2q^3 D} + \frac{1}{2qD}) dV$. The second integral is $\frac{1}{2D} \iiint \frac{dV}{q} = \frac{4\pi R^3/3}{2D^2}$. The first integral over ϕ uses the same $u = D^2 - 2\rho D \cos \phi + \rho^2 = q^2$ as in the text: $\int_0^\pi \frac{\sin \phi d\phi}{q^3} = \int \frac{du/2\rho D}{u^{3/2}} = [\frac{-1}{\rho D u^{1/2}}]_{\phi=0}^{\phi=\pi} = \frac{1}{\rho D} (\frac{1}{D-\rho} - \frac{1}{D+\rho}) = \frac{2}{D(D^2 - \rho^2)}$. The θ integral gives 2π and then the ρ integral is $\int_0^R 2\pi \frac{2}{D(D^2 - \rho^2)} \frac{D^2 - \rho^2}{2D} \rho^2 d\rho = \frac{4\pi R^3/3}{D^2}$. The two integrals give $\frac{4\pi R^3/3}{D^2}$ as Newton hoped and expected.
- 28** The small movement produces a right triangle with hypotenuse ΔD and almost the same angle α . So the new small side Δq is $\Delta D \cos \alpha$.
- 30** $\iint q dA = 4\pi \rho^2 D + \frac{4\pi}{3} \frac{\rho^4}{D}$. Divide by $4\pi \rho^2$ to find $\bar{q} = D + \frac{\rho^2}{3D}$ for the shell. Then the integral over ρ gives $\iiint q dV = \frac{4\pi}{3} R^3 D + \frac{4\pi}{15} \frac{R^5}{D}$. Divide by the volume $\frac{4\pi}{3} R^3$ to find $\bar{q} = D + \frac{R^2}{5D}$ for the solid ball.
- 32** Yes. First concentrate the Earth to a point at its center – this is OK for each point in the Sun. Then concentrate the Sun at its center – this does not change the force on the (concentrated) Earth.
- 34** $J = aci + bfg + cdh - ceg - afh - bdi$.
- 36** Column 1: $\sqrt{\sin^2 \phi (\cos^2 \theta + \sin^2 \theta) + \cos^2 \phi} = 1$; Column 2: $\sqrt{\rho^2 \cos^2 \phi (\cos^2 \theta + \sin^2 \theta) + \rho^2 \sin^2 \phi} = \rho$; Column 3: $\sqrt{\rho^2 \sin^2 \phi (\sin^2 \theta + \cos^2 \theta) + \rho^2 \cos^2 \phi} = \rho \sin \phi$. These are the edge lengths of the box. The dot products of these columns are zero; so $J = \text{volume of box} = (1)(\rho)(\rho \sin \phi)$ as before.
- 38** Column 1: $\sqrt{\cos^2 \theta + \sin^2 \theta} = 1$; Column 2: $\sqrt{r^2 \sin^2 \theta + r^2 \cos^2 \theta} = r$; Column 3: $\sqrt{0^2 + 0^2 + 1^2} = 1$. Again the dot products of the columns are zero and $J = \text{volume of box} = (1)(r)(1) = r$.
- 40** $I = \frac{8}{15}\pi R^5$; $J = \frac{2}{5}$; the mass is closer to the axis.
- 42** The ball comes to a stop at Australia and returns to its starting point. It continues to oscillate in harmonic motion $y = R \cos(\sqrt{c/m} t)$.