

18.04 Recitation 8
Vishesh Jain

- 1.1 Show that if $g(z)$ has a simple zero at z_0 , then $1/g(z)$ has a simple pole at z_0 .
- 1.2. Show that $\text{Res}(1/g, z_0) = 1/g'(z_0)$.
- 1.3. Let $f(z) = 1/\sin(z)$. Find all the poles, show that they are simple, and use the previous part to find the residues at these poles.
- 2.1. Let $p(z)$ and $q(z)$ be analytic at $z = z_0$. Assume $p(z_0) \neq 0$ and q has a simple zero at z_0 . Show that $\text{Res}_{z=z_0}(p(z)/q(z)) = p(z_0)/q'(z_0)$.
- 2.2. Let $f(z) = \cot(z)$. Find all the poles, show that they are simple, and use the previous part to find residues at these poles.
3. By using the Taylor series of $\cos(z)$ and $\sin(z)$ around $z = 0$, compute the first few terms of the Laurent expansion of $\cot(z)$ around $z = 0$.
4. Suppose $f(z)$ is analytic in the region A except for a set of isolated singularities. Suppose C is a simple closed curve in A that doesn't go through any of the singularities of f and is oriented counterclockwise.
- 4.1. Suppose that there is only one isolated singularity inside C at the point z_1 . By using the extended version of Cauchy's theorem, show that $\int_C f(z)dz = 2\pi i \text{Res}(f, z_1)$.
- 4.2. Suppose now that there are two isolated singularities inside C at the points z_1 and z_2 . Again, by using the extended version of Cauchy's theorem, show that $\int_C f(z)dz = 2\pi i (\text{Res}(f, z_1) + \text{Res}(f, z_2))$.
- 4.3. Generalize the previous part to show that

$$\int_C f(z)dz = 2\pi i \left(\sum \text{residues of } f \text{ inside } C \right).$$

This is Cauchy's residue theorem.

MIT OpenCourseWare
<https://ocw.mit.edu>

18.04 Complex Variables with Applications
Spring 2018

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.