

On the decay of crossing numbers

Jacob Fox*

Csaba D. Tóth†

Abstract

The crossing number $\text{cr}(G)$ of a graph G is the minimum number of crossings over all drawings of G in the plane. In 1993, Richter and Thomassen [RT93] conjectured that there is a constant c such that every graph G with crossing number k has an edge e such that $\text{cr}(G - e) \geq k - c\sqrt{k}$. They showed only that G always has an edge e with $\text{cr}(G - e) \geq \frac{2}{5}\text{cr}(G) - O(1)$. We prove that for every connected graph G with n vertices, $m \geq 1$ edges, and every edge e of G , we have $\text{cr}(G - e) \geq \text{cr}(G) - 2m + n/2 + 1$. This confirms the Richter-Thomassen conjecture for dense graphs. We also prove that for every fixed $\epsilon > 0$, there is a constant n_0 depending on ϵ such that if G is a graph with $n > n_0$ vertices and $m > n^{1+\epsilon}$ edges, then G has a subgraph G' with at most $(1 - \frac{1}{24\epsilon})m$ edges such that $\text{cr}(G') \geq (\frac{1}{28} - o(1))\text{cr}(G)$.

Key words: crossing number

1 Introduction

The crossing number $\text{cr}(G)$ of a (simple) graph G is the minimum possible number of crossings in any drawing of G in the plane. A famous result of Ajtai et al. [ACNS82] and Leighton [L84] states that if G is a graph with n vertices and $m \geq 4n$ edges, then

$$\text{cr}(G) \geq \frac{m^3}{64n^2}. \quad (1)$$

For graphs with n vertices and $m \geq \frac{103}{16}n$ edges, Pach et al. [PRTT04] improved Inequality (1) by a constant factor to

$$\text{cr}(G) \geq \frac{1024}{31827} \frac{m^3}{n^2}. \quad (2)$$

It is well known that for every positive integer k , there is a graph G and an edge e of G such that $\text{cr}(G) = k$ but $G - e$ is planar. In 1993, Richter and Thomassen [RT93] conjectured that there is a constant c such that for every nonempty graph G with crossing number k , there is an edge e of G such that $\text{cr}(G - e) \geq k - c\sqrt{k}$. They showed only that G always has an edge e with $\text{cr}(G - e) \geq \frac{2}{5}\text{cr}(G) - O(1)$. Salazar [S00] proved that for every graph G with no vertices of degree 3, there is an edge e of G such that $\text{cr}(G - e) \geq \frac{1}{2}\text{cr}(G) - O(1)$.

We prove the following theorem, which, combined with Inequality (2), improves on the Richter-Thomassen result for graphs with n vertices and $m \geq 10.1n$ edges.

Theorem 1 *For every connected graph G with n vertices, $m \geq 1$ edges, and every edge e of G , we have*

$$\text{cr}(G - e) \geq \text{cr}(G) - 2m + \frac{n}{2} + 1.$$

*Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA.

†Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA.

By Inequality (1), Theorem 1 confirms the Richter-Thomassen conjecture for dense graphs, that is, for graphs with $\Omega(n^2)$ vertices. Moreover, Theorem 1 implies that deleting any edge from a dense graph G reduces the crossing number by at most $O(\sqrt{\text{cr}(G)})$.

The following theorem demonstrates that for every graph G that is not too sparse, we can delete a constant fraction of the edges of G such that the crossing number of the remaining subgraph G' is at least a constant fraction of the crossing number of G .

Theorem 2 *For every $\epsilon > 0$, there is a constant n_0 depending on ϵ such that if G is a graph with $n > n_0$ vertices and $m > n^{1+\epsilon}$ edges, then G has a subgraph G' formed by deleting at least $(\frac{\epsilon}{24})m$ edges from G such that $\text{cr}(G') \geq (\frac{1}{28} - o(1))\text{cr}(G)$.*

To prove Theorem 2, we derive in Section 4 a lower bound on the crossing number that improves on Inequality (1) for graphs with highly irregular degree patterns. Theorem 1 is proved in Section 3 and Theorem 2 is proved in Section 2.

2 Drawing edges with the embedding method

We use the embedding method along the lines of Leighton [L83], Richter and Thomassen [RT93], Shahrokhi et al. [SSSV97], and Székely [S04a]. The embedding method generates a planar drawing (embedding) $D(G)$ of a graph G based on a drawing $D(H)$ of a subgraph $H \subset G$. The drawing $D(G)$ respects $D(H)$ on the edges of H and for every edge $e = (v, w) \in G \setminus H$, the drawing of e follows “infinitesimally close” a path between v and w in the drawing $D(H)$. We can distinguish two categories of crossings that involve edges of $G \setminus H$ in the drawing $D(G)$. A *first category crossing* arises infinitesimally close to a crossing in $D(H)$. A *second category crossing* arises infinitesimally close to a vertex in $D(H)$.

Before we prove Theorem 2, we illustrate the embedding method with a bound on the minimum decay of the crossing number after deleting one edge. This improves on the Richter-Thomassen bound for graphs with $m \geq 7.66n$ edges.

Proposition 1 *For every connected graph G with n vertices and m edges, there is an edge e of G such that*

$$\text{cr}(G - e) \geq \frac{p}{p+2} \left(\text{cr}(G) - m + \frac{n}{2} \right),$$

where $p = \lceil \frac{m}{n-1} - 1 \rceil$.

The proof of Proposition 1 follows immediately from Proposition 2 and Lemma 1 below. Nagamochi and Ibaraki [NI92] proved the following lemma, which is a slight variant of Mader’s theorem, and shows that every graph with n vertices and m edges has a pair of adjacent vertices with at least $\frac{m}{n-1}$ edge-disjoint paths between them.

Lemma 1 (Mader, Nagamochi and Ibaraki) *If G is a graph with m edges and n vertices, then there is an edge $e = (v, w)$ of G such that there are at least $\frac{m}{n-1} - 1$ edge-disjoint paths between v and w in $G - e$.*

Proof: Delete maximal spanning forests F_1, F_2, \dots, F_j one after the other until all edges are deleted. If $e = (v, w)$ is an edge of F_j , then there is a path between v and w in F_i for every $i, 1 \leq i \leq j$. Hence, there are at least $j - 1$ edge-disjoint paths between v and w that do not pass through e . Since each F_i is a forest, it has at most $n - 1$ edges, and so we have $m \leq j(n - 1)$. Substituting, there are at least $\frac{m}{n-1} - 1$ edge-disjoint paths between v and w in $G - e$. \square

Proposition 2 Let G be a connected graph with n vertices and m edges, and $e = (v, w)$ be an edge of G such that there are $p \geq 1$ edge-disjoint paths between v and w in $G - e$. Then

$$\text{cr}(G) \leq \left(1 + \frac{2}{p}\right) \text{cr}(G - e) + m - \frac{n}{2}.$$

Proof: Let D be a drawing of $G - e$ in the plane with $\text{cr}(G - e)$ crossings. Let P_1, P_2, \dots, P_p be p edge-disjoint paths between v and w . Consider the drawing D_j of G in the plane that respects the drawing D of $G - e$ and the edge e follows infinitesimally close to the path P_j between v and w with all loops (and self-crossings) deleted. Let k_j be the number of first category crossings in D_j . Since the paths P_1, P_2, \dots, P_p are edge-disjoint, the drawings D_1, D_2, \dots, D_p of G jointly have at most two first category crossings at each crossing of D : at most two crossings between edges of $G - e$ and different drawings of e , as depicted in Figure1(a). Hence,

$$\sum_{j=1}^p k_j \leq 2\text{cr}(G - e).$$

Therefore, there is an index j , $1 \leq j \leq p$, such that $k_j \leq 2\text{cr}(G - e)/p$.




Figure 1: Drawings of edge e along two edge-disjoint path P_j and $P_{j'}$ may give two first category crossings at a crossing of D (a). If a path P_j traverses a vertex u (b), then the edge e drawn along P_j can choose between two possible routes around u (c–d).

At each internal vertex u of a path P_j , the drawing of e in D_j can take two possible routes, as depicted in Figure 1 (c–d). The two possible routes have a total of $\deg(u) - 1$ second category crossings at u . We draw e along the route with fewer second category crossings, and so there are at most $\frac{1}{2}(\deg(u) - 1)$ crossing at vertex u . Hence, the total number of second category crossings is at most $m - \frac{n}{2}$. Therefore, in the drawing D_j of G , there are at most $(1 + \frac{2}{p})\text{cr}(G - e) + m - \frac{n}{2}$ crossings. \square

The following theorem establishes Theorem 2 for all graphs with n vertices of degree d_1, \dots, d_n such that $\text{cr}(G) \geq \frac{7}{16} \sum_{i=1}^n d_i^2$. For graphs that do not satisfy this condition, we alter the proof utilizing Lemma 3 in Section 4.

Theorem 3 For every ϵ , $0 < \epsilon < 1$, there is a positive constant $n(\epsilon)$ such that for every G with $n > n(\epsilon)$ vertices, with a degree sequence d_1, \dots, d_n , and $m > n^{1+\epsilon}$ edges, there is a subgraph G' of G with at most $(1 - \frac{\epsilon}{8})m$ edges such that

$$4\text{cr}(G') \geq \text{cr}(G) - \frac{3}{8} \sum_{i=1}^n d_i^2.$$

Proof: Erdős and Simonovits [ES82] proved that for every integer $r > 1$, there is a constant c_r such that every graph G with n vertices and $m > c_r n^{1+\frac{1}{r}}$ edges contains a cycle of length $2r$. This implies that for $0 < \epsilon < 1$, there is a positive integer r satisfying $\frac{1}{\epsilon} < r \leq \frac{2}{\epsilon}$ so that every sufficiently large graph G with $m > n^{1+\epsilon}$ edges contains a family \mathcal{C} of edge-disjoint cycles of length $2r$ that cover at least half of the edges of G . Let G' be a subgraph of G formed by deleting an arbitrary edge e_j from each cycle $C_j \in \mathcal{C}$. The remaining edges of cycle C_j form a path P_j . Hence, the number of edges of $G \setminus G'$ is at least $\frac{\epsilon}{8}m$. Let us denote the vertices of G by $v_i, i = 1, 2, \dots, n$, such that the degree of v_i is d_i in G and d'_i in G' . Let $h_i = d_i - d'_i$, which is the number of edges incident to v_i in $G \setminus G'$.

Consider a drawing D' of G' in the plane with $\text{cr}(G')$ crossings. We generate a drawing D of G based on D' by applying the embedding method. In particular, for every edge e_j of cycle $C_j \in \mathcal{C}$, we draw e_j along the path P_j . Since the paths $P_j, j \in \mathcal{C}$ are edge-disjoint, D has at most 4 crossings at every crossing of D' . The total number of crossings of D' and first category crossings of D is at most $4\text{cr}(G')$.

Next we estimate the number of second category crossings. Each of the h_i edges incident to v_i in $G \setminus G'$ is drawn, in a neighborhood of v_i , close to one of the d'_i edges incident to v_i in G' . The vertex v_i with degree d'_i in G' is an internal node of at most $\lfloor (d'_i - h_i)/2 \rfloor$ paths P_j . For every such path P_j , the edge e_j is drawn along one of two possible routes, as depicted in Figure 1(c-d), with the minimum number of crossings with the edges of G incident to the vertex v_i . Every edge $e_j \in G \setminus G'$ passing through a small neighborhood of v_i has at most $\lfloor (d'_i + h_i - 1)/2 \rfloor$ second category crossings with edges of G incident to v_i . Each pair of edges passing through a small neighborhood of v_i cross at most once. So the total number of second category crossings at v_i is at most

$$\left\lfloor \frac{d'_i - h_i}{2} \right\rfloor \cdot \left\lfloor \frac{d'_i + h_i - 1}{2} \right\rfloor + \binom{\lfloor (d'_i - h_i)/2 \rfloor}{2} < \frac{3}{8}d_i'^2 \leq \frac{3}{8}d_i^2.$$

Summing over all vertices, we have at most $\sum_{i=1}^n \frac{3}{8}d_i^2$ second category crossings.

Hence, we have

$$\text{cr}(G) \leq 4\text{cr}(G') + \frac{3}{8} \sum_{i=1}^n d_i^2.$$

□

We now prove Theorem 2 in general. Order the vertices v_1, v_2, \dots, v_n of G such that their degree sequence d_1, d_2, \dots, d_n monotone increases. Let ℓ be the integer such that $\sum_{i=1}^{\ell-1} d_i < \frac{4m}{3}$ but $\sum_{i=1}^{\ell} d_i \geq \frac{4m}{3}$. Consider the graph G_0 induced by the vertices $v_1, v_2, \dots, v_{\ell}$. Notice that G_0 has at least $\frac{m}{3}$ edges. We choose a family \mathcal{C} of edge-disjoint cycles of length at most $\frac{4}{\epsilon}$ from G_0 so that at least half of the edges of G_0 are covered by cycles of \mathcal{C} . Let G' be a subgraph of G formed by deleting an edge e_j from each cycle $C_j \in \mathcal{C}$. We have deleted at least $\frac{1}{2} \cdot \frac{\epsilon}{4} \cdot \frac{m}{3} = \frac{\epsilon}{24}m$ edges. Let m' be the number of edges of G' and d'_i be the degree of v_i in G' . We have $d'_i \leq d_i$ for $1 \leq i \leq \ell$ and $d'_i = d_i$ for $i > \ell$. It follows that $\sum_{i=1}^{\ell-1} d'_i < \frac{4m'}{3}$. By Lemma 3 (in Section 4 below), we have $\text{cr}(G') \geq (\frac{1}{64} - o(1)) \sum_{i=1}^{\ell} d_i'^2$. If we apply the embedding method to draw graph G based on the drawing of G' with $\text{cr}(G')$ crossings and drawing each e_j along P_j , we obtain

$$\text{cr}(G) \leq 4\text{cr}(G') + \frac{3}{8} \sum_{i=1}^{\ell} d_i'^2.$$

Hence, we have $\text{cr}(G) \leq 4\text{cr}(G') + \frac{3}{8}(64 + o(1))\text{cr}(G') = (28 + o(1))\text{cr}(G')$.

3 Deleting one edge

In this section, we prove Theorem 1. Suppose G is a simple connected graph, $e = (v, w)$ is an edge of $G = (V, E)$, and D is a drawing of $G - e$ with $\text{cr}(G - e)$ crossings. We may further suppose that v and w are in the same component of $G - e$ since otherwise $\text{cr}(G - e) = \text{cr}(G)$. Two components can be drawn in two disjoint halfplanes such that v and w are on the outer face in either; then v and w are in the same face of the union and so one can add the edge vw without crossings. Let C denote the set of crossings in the drawing D . We define a planar graph $H = (W, F)$ on the drawing of $G - e$. The vertex set of H is $W = V \cup C$ and two vertices are connected by an edge if they are consecutive vertices along (the drawing of) an edge of $G - e$. An example, where G is the Petersen graph, is depicted in Figure 2




Figure 2: Two drawings of the Petersen graph (a–b) and the planar graph corresponding to the second drawing (c).

For each vertex $v \in W$, assign a weight $w(v)$ defined by

$$w(v) = \begin{cases} \deg_{G-e}(v) & \text{if } v \text{ is a vertex of } G - e \\ 1 & \text{if } v \text{ is a crossing of } G - e \end{cases} \quad (3)$$

For every path $P = (p_1, p_2, \dots, p_k)$ in H , we define the weight $w(P)$ to be the sum of the weights of its internal vertices $w(P) = \sum_{i=2}^{k-1} w(p_i)$. Let P_e be a minimum weight path between v and w . We may assume that P_e is a simple path, since removing loops strictly decreases the weight. We generate a drawing D' of G based on the drawing D of $G - e$ by adding the edge e infinitesimally close to the path P_e .




Figure 3: A path approaching a crossing can traverse it in three different ways depending on its destination.

In Figure 3, we specify how the drawing of e traverses a crossing in D along P_e . At each crossing of D that P_e traverses, e has at most one new first category crossing. We now show that e has at most one first category crossing with every edge of $G - e$. Suppose that e and an edge $e' \in G - e$

have more than one first category crossings in D' . This means that the path P_e has at least two crossings with e' in D . Consider two consecutive crossings of P_e and e' . There are two ways as two simple curves can cross twice, as depicted in Figure 4(a-b). Since P_e has minimal weight, we can redraw e' so as to have a drawing of $G - e$ with fewer crossings than in D , contradicting that D has $\text{cr}(G - e)$ crossings. This redrawing of e' is depicted in Figure 4(c-d).




Figure 4: Two possible configuration of two consecutive crossings of e and e' (a-b). Redrawing of e' along the drawing of e (c-d).

We have shown that our drawing of e has at most $m - 1$ first category crossings with edges of $G - e$. We have yet to specify how edge e is drawn around vertices of $G - e$ along P_e . There are two possible routes that P_e might take at each vertex u of $G - e$, as shown in Figure 1(b-d). Pick the route with fewer crossings. The two possible routes jointly have a total of $\deg_{G-e}(u) - 1$ crossings, so the route with fewer crossings has at most $\frac{1}{2}(\deg_{G-e}(u) - 1) \leq \frac{1}{2}(\deg_G(u) - 1)$ second category crossings at u . Summing over all vertices of G , we have at most $m - \frac{n}{2}$ second category crossings. The sum of the number of crossings in D , the number of first and second category crossings in the drawing D' is at most $\text{cr}(G - e) + 2m - \frac{n}{2} - 1$, establishing Theorem 1. \square

4 The sum of degree squares and the crossing number

The *bisection width*, denoted by $b(G)$, is defined for every simple graph G with at least two vertices. $b(G)$ is the smallest nonnegative integer such that there is a partition of the vertex set $V = V_1 \cup^* V_2$ with $\frac{1}{3} \cdot |V| \leq |V_i| \leq \frac{2}{3} \cdot |V|$ for $i = 1, 2$, and $|E(V_1, V_2)| \leq b(G)$. Extending the Lipton-Tarjan separator theorem [LT79], Gazit and Miller [GM90] established an upper bound on the bisection width in terms of the sum of degree squares.

Theorem 4 (Gazit and Miller) *Let G be a planar graph with n vertices of degree d_1, d_2, \dots, d_n . Then*

$$b^2(G) \leq \frac{5 + 2\sqrt{6}}{4} \cdot \sum_{i=1}^n d_i^2.$$

Pach, Shahrokhi, and Szegedy [PSS96] used Theorem 4 to relate the bisection width with the crossing number.

Theorem 5 (Pach, Shahrokhi, and Szegedy) *Let G be a graph with n vertices of degree d_1, d_2, \dots, d_n . Then*

$$40\text{cr}(G) \geq b^2(G) - \frac{5}{2} \cdot \sum_{i=1}^n d_i^2(G).$$

Pach, Spencer and Tóth [PST00] have further exploited the connection between the bisection width and the crossing number. They have established lower bounds on the crossing number of graphs with some monotone graph property in terms of the number of edges and vertices of the graph. A simplified version of their proof method yields the following bounds.

Lemma 2 *Let $G(V, E)$ be a graph with n vertices of degree d_1, d_2, \dots, d_n , and m edges. For any δ , $0 < \delta < 1$, let $\Delta = \Delta(\delta)$ be the integer such that $\sum_{i=1}^n \min(d_i, \Delta) < 2\delta m$ but $\sum_{i=1}^n \min(d_i, \Delta + 1) \geq 2\delta m$. The crossing number of G is bounded by the sum of truncated degree squares. If $m \geq 45(1 - \delta)^{-2}n \log^2 n$, then*

$$\text{cr}(G) \geq \frac{1}{16} \sum_{i=1}^n (\min(d_i, \Delta))^2.$$

Lemma 3 *Let $G(V, E)$ be a graph with n vertices and m edges, and let $d_1 \leq d_2 \leq \dots \leq d_n$ denote the degree sequence sorted in monotone increasing order. Let ℓ be the integer such that $\sum_{i=1}^{\ell-1} d_i < 4m/3$ but $\sum_{i=1}^{\ell} d_i \geq 4m/3$. The crossing number of G is bounded by a prefix sum of the degree squares. If $m = \Omega(n \log^2 n)$, then*

$$\text{cr}(G) \geq \left(\frac{1}{64} - o(1) \right) \sum_{i=1}^{\ell} d_i^2.$$

These bounds are better than the classical lower bound (1) due to Ajtai et al. [ACNS82] and Leighton [L84] for graphs of irregular degree patterns and $m = O(n^{3/2})$ edges. Consider the complete bipartite graph $K_{a,b}$ with $n = a + b$ vertices and $m = ab$ edges, where $a \leq b$. For this graph, our Lemma 3 gives $\text{cr}(G) = \Omega(ab^2)$, which is a tighter lower bound than the classical $\Omega(m^3/n^2) = \Omega(a^3b)$ for $a \leq b < a^2$. Similar bounds have also been deduced by Pach and Solymosi [PS05].

Proof of Lemma 2. We decompose the graph G by the following recursive algorithm into induced subgraphs such that every subgraph is either a singleton or its squared bisection width is at least five times the sum of its degree squares. In an induced subgraph $H \subseteq G$, we denote by $\deg_H(v)$ the degree of a vertex $v \in V(H)$.

1. Let $S_0 = \{G\}$ and $i = 0$.
2. Repeat until $|V(H)| = 1$ or $b^2(H) \geq 5 \sum_{v \in H} \deg_H^2(v)$ for every $H \in S_i$.

Set $i := i + 1$ and $S_{i+1} := \emptyset$. For every $H \in S_i$, do

- If $b^2(H) \geq 5 \sum_{v \in H} \deg_H^2(v)$ or $|V(H)| \leq (2/3)^i |V|$, then let $S_{i+1} := S_{i+1} \cup \{H\}$;
- otherwise split H into graphs H_1 and H_2 along an edge separator of size $b(H)$, and let $S_{i+1} := S_{i+1} \cup \{H_1, H_2\}$.

3. Return S_i .

First, we show that the algorithm is correct. In every round, every graph $H \in S_i$ that does not satisfy the end condition has at most $|V(H)| \leq (2/3)^i \cdot |V|$ vertices. The algorithm terminates in $t \leq \log_{(3/2)} n$ rounds, and it returns a set S_t of induced subgraphs. By Theorem 5 and the end condition of the decomposition algorithm, for every $H \in S_t$ we have $40\text{cr}(H) \geq (5/2) \sum_{v \in H} \deg_H^2(v)$. So

$$40\text{cr}(G) \geq 40 \sum_{H \in S_t} \text{cr}(H) \geq \frac{5}{2} \cdot \sum_{H \in S_t} \sum_{v \in H} \deg_H^2(v) \geq \frac{5}{2} \cdot \sum_{v \in V} \deg_{H(v,t)}^2(v), \quad (4)$$

where $H(v, i)$ denotes the graph $H \in S_i$ containing vertex $v \in V$.

Next, we count the number of edges deleted during the recursive decomposition. Following an argument of [PST00], we count separately the edges deleted in each step of the decomposition algorithm. Let $S'_i = \{H : H \in S_i, H \notin S_{i+1}\}$, that is, S'_i consists of those subgraphs in S_i that are decomposed at step i . Notice that $|S'_i| < (\frac{3}{2})^{i+1}$ since every subgraph of S_i that splits has more than $(2/3)^{i+1}|V|$ vertices. Let $V_i = \{v : v \text{ is a vertex of a graph } H \in S'_i\}$.

In step i , when some of the subgraphs in S_i are decomposed in S_{i+1} , the total number of deleted edges is at most

$$\sum_{H \in S'_i} \sqrt{5 \sum_{v \in H} \deg_H^2(v)}.$$

Using the Cauchy-Schwartz inequality, we have

$$\sum_{H \in S'_i} \sqrt{5 \sum_{v \in H} \deg_H^2(v)} \leq \sqrt{5|S'_i|} \sqrt{\sum_{v \in V_i} \deg_{H(v,i)}^2(v)} \leq \sqrt{5 \left(\frac{3}{2}\right)^{i+1}} \sqrt{\sum_{v \in V_i} \deg_{H(v,i)}^2(v)}.$$

Since $|V(H)| \leq (\frac{2}{3})^i |V|$ for each subgraph $H \in S'_i$, we conclude that

$$\begin{aligned} \sqrt{5 \left(\frac{3}{2}\right)^{i+1}} \sqrt{\sum_{v \in V_i} \deg_{H(v,i)}^2(v)} &\leq \sqrt{5 \left(\frac{3}{2}\right)^{i+1}} \sqrt{\max_{v \in V_i} \deg_{H(v,i)}(v) \cdot \sum_{v \in V_i} \deg_{H(v,i)}(v)} \\ &\leq \sqrt{5 \left(\frac{3}{2}\right)^{i+1}} \sqrt{\left(\frac{2}{3}\right)^i n(2m)} \leq \sqrt{15mn}. \end{aligned}$$

Since the algorithm terminates in at most $\log n / \log(3/2)$ steps, the total number of edges deleted throughout the decomposition algorithm is at most

$$\frac{\sqrt{15}}{\log(3/2)} \sqrt{mn} \log n.$$

This is less than $(1 - \delta)m$ if $m \geq \frac{15}{\log^2(3/2)} (1 - \delta)^{-2} n \log^2 n$, or if $m \geq 45(1 - \delta)^{-2} n \log^2 n$.

We are now ready to estimate $\sum_{v \in V} \deg_{H(v,t)}^2(v)$. Since the number of edges decreased by at most $(1 - \delta)m$, the sum of degrees decreased by at most $(2 - 2\delta)m$. The sum of degree squares decreases maximally if the highest degrees are truncated to at most Δ , and so we have

$$\sum_{v \in V} \deg_{H(v,t)}^2(v) \geq \sum_{i=1}^n (\min(d_i, \Delta))^2. \quad (5)$$

This completes the proof of Lemma 2. \square




Figure 5: The monotone increasing degree sequence of a graph G .

Proof of Lemma 3. We extend the argument of the previous proof with $\delta = \frac{5}{6}$. If $d_\ell \leq \Delta$, then the right hand side of (5) must clearly be at least $\sum_{i=1}^\ell d_i^2$ and our proof is complete. Let us assume that $\Delta < d_\ell$. Refer to Figure 5.

Recall that $\sum_{i=1}^n d_i = 2m$. We have assumed that $\sum_{i=\ell+1}^n d_i \leq \frac{2m}{3} < \sum_{i=\ell}^n d_i$, and for $\delta = \frac{5}{6}$ we have $\sum_{i=1}^n \min(d_i, \Delta) < \frac{5m}{3} \leq \sum_{i=1}^n \min(d_i, \Delta + 1)$. It follows that $(n - \ell + 1)(\Delta + 1) > \frac{m}{3}$. Since $\Delta < n$ and $n = o(m)$, we conclude that $(n - \ell)\Delta > (1 - o(1))\frac{m}{3}$.

Observe that $(n - \ell)d_\ell \leq \sum_{i=\ell+1}^n d_i \leq \frac{2m}{3}$, and so $m \geq \frac{3}{2}(n - \ell)d_\ell$. Furthermore, observe that $\sum_{i=1}^\ell \max(0, d_i - \Delta) \leq \sum_{i=1}^n \max(0, d_i - \Delta) \leq n + \sum_{i=1}^n \max(0, d_i - (\Delta + 1)) \leq n + \frac{m}{3} = (1 + o(1))\frac{m}{3}$. Putting these simple observations together, we obtain

$$\begin{aligned}
\sum_{i=\ell+1}^n (\min(d_i, \Delta))^2 &= (n - \ell)\Delta^2 > \left(\frac{1}{3} - o(1)\right) m\Delta \geq \left(\frac{1}{2} - o(1)\right) (n - \ell)d_\ell\Delta \\
&\geq \left(\frac{1}{6} - o(1)\right) d_\ell m \geq \left(\frac{1}{2} - o(1)\right) d_\ell \sum_{i=1}^\ell \max(0, d_i - \Delta) \\
&\geq \left(\frac{1}{2} - o(1)\right) \sum_{i=1}^\ell (\max(0, d_i - \Delta))^2.
\end{aligned}$$

We can now estimate the right hand side of Inequality (5).

$$\begin{aligned}
\sum_{i=1}^n (\min(d_i, \Delta))^2 &= \sum_{i=1}^\ell (\min(d_i, \Delta))^2 + \sum_{i=\ell+1}^n (\min(d_i, \Delta))^2 \\
&\geq \sum_{i=1}^\ell (\min(d_i, \Delta))^2 + \left(\frac{1}{2} - o(1)\right) \sum_{i=1}^\ell (\max(0, d_i - \Delta))^2 \\
&\geq \left(\frac{1}{2} - o(1)\right) \sum_{i=1}^\ell (\min(d_i, \Delta))^2 + (\max(0, d_i - \Delta))^2 \geq \left(\frac{1}{4} - o(1)\right) \sum_{i=1}^\ell d_i^2.
\end{aligned}$$

Comparing the above inequality with Inequalities (4) and (5), we obtain $\text{cr}(G) \geq (\frac{1}{64} - o(1)) \sum_{i=1}^\ell d_i^2$.
 \square

5 Acknowledgments

We would like to thank Daniel J. Kleitman, János Pach, and Rados Radoičić for helpful comments. Thanks to László Szegő for directing us to Nagamochi and Ibaraki's results.

References

- [ACNS82] M. Ajtai, V. Chvátal, M. Newborn, and E. Szemerédi: Crossing-free subgraphs, in *Theory and Practice of Combinatorics*, vol. 60 of Mathematical Studies, North-Holland, Amsterdam, 1982, pp. 9–12.
- [ES82] P. Erdős and M. Simonovits: Compactness results in extremal graph theory, *Combinatorica* **2** (1982), 275–288.
- [GM90] H. Gazit and G. L. Miller: Planar separators and the Euclidean norm, *SIGAL International Symposium on Algorithms*, vol. 450 of LNCS, Springer-Verlag, 1990, pp. 338–347.
- [L83] T. Leighton: *Complexity Issues in VLSI*, MIT Press, Cambridge, MA, 1983,
- [L84] T. Leighton: New lower bound techniques for VLSI, *Math. Systems Theory* **17** (1984), 47–70.
- [LT79] R. J. Lipton and R. E. Tarjan: A separator theorem for planar graphs, *SIAM J. Appl. Math.* **36** (1979), 177–189.
- [NI92] H. Nagamochi and T. Ibaraki: A linear-time algorithm for finding a sparsek-connected spanning subgraph of ak-connected graph, *Algorithmica* **7** (1992), 583–596.
- [PRTT04] J. Pach, R. Radoičić, G. Tardos, and G. Tóth: Improving the Crossing Lemma by finding more crossings in sparse graphs, *20th ACM Symposium on Computational Geometry*, ACM Press, New York, 2004, 68–75
- [PSS96] J. Pach, F. Shahrokhi, and M. Szegedy: Applications of the crossing number, *Algorithmica* **16** (1996), 111–117.
- [PS05] J. Pach and J. Solymosi: personal communication, 2005.
- [PST00] J. Pach, J. Spencer, and G. Tóth: New bounds on crossing numbers, *Discrete Comput. Geom.* **24** (2000), no. 4, 623–644.
- [RT93] B. Richter and C. Thomassen: Minimal graphs with crossing number at least k , *J. Combin. Theory Ser. B* **58** (1993), 217–224.
- [S00] G. Salazar: On a crossing number result of Richter and Thomassen, *J. Combin. Theory Ser. B* **79** (2000), 98–99.
- [S04] L. A. Székely: A successful concept for measuring non-planarity of graphs: the crossing number. *Discrete Math.* **276** (2004), no. 1-3, 331–352.
- [S04a] L. A. Székely: Short proof for a theorem of Pach, Spencer, and Tóth. *Towards a theory of geometric graphs*, 281–283, Contemp. Math., 342, Amer. Math. Soc., Providence, RI, 2004.
- [SSSV97] F. Shahrokhi, O. Sýkora, L. A. Székely, and I. Vrto: Crossing numbers: bounds and applications. *Intuitive geometry* (Budapest, 1995), 179–206, Bolyai Soc. Math. Stud., 6, János Bolyai Math. Soc., Budapest, 1997.