

### 13. The Poincaré-Birkhoff-Witt theorem

#### 13.1. The statement of the Poincaré-Birkhoff-Witt theorem.

Let  $\mathfrak{g}$  be a Lie algebra. Define a filtration<sup>10</sup> on  $U(\mathfrak{g})$  by setting  $\deg(\mathfrak{g}) = 1$ . Thus  $F_n U(\mathfrak{g})$  is the image of  $\bigoplus_{i=0}^n \mathfrak{g}^{\otimes i} \subset T\mathfrak{g}$ . Note that since

$$xy - yx = [x, y], \quad x \in \mathfrak{g},$$

we have  $[F_i U(\mathfrak{g}), F_j U(\mathfrak{g})] \subset F_{i+j-1} U(\mathfrak{g})$ . Thus,  $\text{gr}U(\mathfrak{g})$  is commutative; in other words, we have a surjective algebra morphism  $\phi : S\mathfrak{g} \rightarrow \text{gr}U(\mathfrak{g})$ .

**Theorem 13.1.** (*Poincaré-Birkhoff-Witt theorem*) *The homomorphism  $\phi$  is an isomorphism.*

We will prove Theorem 13.1 in Subsection 13.2. Now let us discuss its reformulation in terms of a basis and corollaries.

Given a basis  $\{x_i\}$  of  $\mathfrak{g}$ , fix an ordering on this basis and consider ordered monomials  $\prod_i x_i^{n_i}$ , where the product is ordered according to the ordering of the basis. The statement that  $\phi$  is surjective is equivalent to saying that ordered monomials span  $U(\mathfrak{g})$ . This is also easy to see directly: any monomial can be ordered using the commutation relations at the cost of an error of lower degree, so proceeding recursively, we can write any monomial as a linear combination of ordered ones. Thus the PBW theorem can be formulated as follows:

**Theorem 13.2.** *The ordered monomials are linearly independent, hence form a basis of  $U(\mathfrak{g})$ .*

**Corollary 13.3.** *The map  $\rho : \mathfrak{g} \rightarrow U(\mathfrak{g})$  is injective. Thus  $\mathfrak{g} \subset U(\mathfrak{g})$ .*

**Remark 13.4.** Let  $\mathfrak{g}$  be a vector space equipped with a bilinear map  $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ . Then one can define the algebra  $U(\mathfrak{g})$  as above. However, if the map  $\rho : \mathfrak{g} \rightarrow U(\mathfrak{g})$  is injective then we clearly must have  $[x, x] = 0$  for  $x \in \mathfrak{g}$  and the Jacobi identity, i.e.,  $\mathfrak{g}$  has to be a Lie algebra. Thus the PBW theorem and even Corollary 13.3 fail without the axioms of a Lie algebra.

**Corollary 13.5.** *Let  $\mathfrak{g}_i$ ,  $1 \leq i \leq n$ , be Lie subalgebras of  $\mathfrak{g}$  such that  $\mathfrak{g} = \bigoplus_i \mathfrak{g}_i$  as a vector space (but  $[\mathfrak{g}_i, \mathfrak{g}_j]$  need not be zero). Then the multiplication map  $\bigotimes_i U(\mathfrak{g}_i) \rightarrow U(\mathfrak{g})$  in any order is a linear isomorphism.*

*Proof.* The corollary follows immediately from the PBW theorem by choosing a basis of each  $\mathfrak{g}_i$ .  $\square$

<sup>10</sup>The grading on  $T\mathfrak{g}$  does not descend to  $U(\mathfrak{g})$ , in general, since the relation  $xy - yx = [x, y]$  is not homogeneous: the right hand side has degree 1 while the left hand side has degree 2. So  $U(\mathfrak{g})$  is not graded but is only filtered.

**Remark 13.6.** 1. Corollary 13.5 applies to the case of infinitely many  $\mathfrak{g}_i$  if we understand the tensor product accordingly: the span of tensor products of elements of  $U(\mathfrak{g}_i)$  where almost all of these elements are equal to 1.

2. Note that if  $\dim \mathfrak{g}_i = 1$ , this recovers the PBW theorem itself, so Corollary 13.5 is in fact a generalization of the PBW theorem.

Let  $\text{char}(\mathbf{k}) = 0$ . Define the **symmetrization map**  $\sigma : S\mathfrak{g} \rightarrow U(\mathfrak{g})$  given by

$$\sigma(y_1 \otimes \dots \otimes y_n) = \frac{1}{n!} \sum_{s \in S_n} y_{s(1)} \dots y_{s(n)}.$$

It is easy to see that this map commutes with the adjoint action of  $\mathfrak{g}$ .

**Corollary 13.7.**  $\sigma$  is an isomorphism.

*Proof.* It is easy to see that  $\text{gr}\sigma$  (the induced map on the associated graded algebra) coincides with  $\phi$ , so the result follows from the PBW theorem.  $\square$

Let  $Z(U(\mathfrak{g}))$  denote the center of  $U(\mathfrak{g})$ .

**Corollary 13.8.** The map  $\sigma$  defines a filtered vector space isomorphism  $\sigma_0 : Z(U(\mathfrak{g})) \rightarrow (S\mathfrak{g})^{\text{adg}}$  whose associated graded is the algebra isomorphism  $\phi|_{Z(U(\mathfrak{g}))} : Z(U(\mathfrak{g})) \rightarrow (S\mathfrak{g})^{\text{adg}}$ .

In the case when  $\mathfrak{g} = \text{Lie}G$  for a connected Lie group  $G$ , we thus obtain a filtered vector space isomorphism of the center of  $U(\mathfrak{g})$  with  $(S\mathfrak{g})^{\text{Ad}G}$ .

**Remark 13.9.** The map  $\sigma_0$  is not, in general, an algebra homomorphism; however, a nontrivial theorem of M. Duflo says that if  $\mathfrak{g}$  is finite dimensional then there exists a canonical filtered *algebra isomorphism*  $\eta : Z(U(\mathfrak{g})) \rightarrow (S\mathfrak{g})^{\text{adg}}$  (a certain twisted version of  $\sigma_0$ ) whose associated graded is  $\phi|_{Z(U(\mathfrak{g}))}$ . A construction of the Duflo isomorphism can be found in [CR].

**Example 13.10.** Let  $\mathfrak{g} = \mathfrak{sl}_2 = \mathfrak{so}_3$ . Then  $\mathfrak{g}$  has a basis  $x, y, z$  with  $[x, y] = z$ ,  $[y, z] = x$ ,  $[z, x] = y$ , and  $G = SO(3)$  acts on these elements by ordinary rotations of the 3-dimensional space. So the only  $G$ -invariant polynomials of  $x, y, z$  are polynomials of  $r^2 = x^2 + y^2 + z^2$ . Thus we get that  $Z(U(\mathfrak{g})) = \mathbb{C}[x^2 + y^2 + z^2]$ . In terms of  $e, f, h$ , we have

$$x^2 + y^2 + z^2 = -4fe - h^2 - 2h = -2C,$$

where  $C$  is the Casimir element.

**13.2. Proof of the PBW theorem.** The proof of Theorem 13.1 is based on the following key lemma.

**Lemma 13.11.** *There exists a unique linear map  $\varphi : T\mathfrak{g} \rightarrow S\mathfrak{g}$  such that*

(i) *for an ordered monomial  $X := x_{i_1} \dots x_{i_m} \in \mathfrak{g}^{\otimes m}$  one has  $\varphi(X) = X$ ;*

(ii) *one has  $\varphi(I) = 0$ ; in other words,  $\varphi$  descends to a linear map  $\bar{\varphi} : U(\mathfrak{g}) \rightarrow S\mathfrak{g}$ .*

**Remark 13.12.** The map  $\varphi$  is not canonical and depends on the choice of the ordered basis  $x_i$  of  $\mathfrak{g}$ .

Note that Lemma 13.11 immediately implies the PBW theorem, since by this lemma the images of ordered monomials under  $\varphi$  are linearly independent in  $S\mathfrak{g}$ , implying that these monomials themselves are linearly independent in  $U(\mathfrak{g})$ .

*Proof.* It is clear that  $\varphi$  is unique if it exists since ordered monomials span  $U(\mathfrak{g})$ . We will construct  $\varphi$  by defining it inductively on  $F_n T\mathfrak{g}$  for  $n \geq 0$ .

Suppose  $\varphi$  is already defined on  $F_{n-1} T\mathfrak{g}$  and let us extend it to  $F_n T\mathfrak{g} = F_{n-1} T\mathfrak{g} \oplus \mathfrak{g}^{\otimes n}$ . So we should define  $\varphi$  on  $\mathfrak{g}^{\otimes n}$ . Since  $\varphi$  is already defined on ordered monomials  $X$  (by  $\varphi(X) = X$ ), we need to extend this definition to all monomials.

Namely, let  $X$  be an ordered monomial of degree  $n$ , and let us define  $\varphi$  on monomials of the form  $s(X)$  for  $s \in S_n$ , where

$$s(y_1 \dots y_n) := y_{s(1)} \dots y_{s(n)}.$$

To this end, fix a decomposition  $D$  of  $s$  into a product of transpositions of neighbors:

$$s = s_{j_r} \dots s_{j_1},$$

and define  $\varphi(s(X))$  by the formula

$$\varphi(s(X)) := X + \Phi_D(s, X),$$

where

$$\Phi_D(s, X) := \sum_{m=0}^{r-1} \varphi([\cdot, \cdot]_{j_{m+1}}(s_{j_m} \dots s_{j_1}(X))),$$

and

$$[\cdot, \cdot]_j(y_1 \dots y_j y_{j+1} \dots y_n) := y_1 \dots [y_j, y_{j+1}] \dots y_n.$$

We need to show that  $\varphi(s(X))$  is well defined, i.e.,  $\Phi_D(s, X)$  does not really depend on the choice of  $D$  and  $s$  but only on  $s(X)$ . We first show that  $\Phi_D(s, X)$  is independent on  $D$ .

To this end, recall that the symmetric group  $S_n$  is generated by  $s_j, 1 \leq j \leq n-1$  with defining relations

$$s_j^2 = 1; s_j s_k = s_k s_j, |j - k| \geq 2; s_j s_{j+1} s_j = s_{j+1} s_j s_{j+1}.$$

Thus any two decompositions of  $s$  into a product of transpositions of neighbors can be related by a sequence of applications of these relations somewhere inside the decomposition.

Now, the first relation does not change the outcome by the identity  $[x, y] = -[y, x]$ .

For the second relation, suppose that  $j < k$  and we have two decompositions  $D_1, D_2$  of  $s$  given by  $s = ps_j s_k q$  and  $s = ps_k s_j q$ , where  $q$  is a product of  $m$  transpositions of neighbors. Let  $q(X) = YabZcdT$  where  $a, b, c, d \in \mathfrak{g}$  stand in positions  $j, j+1, k, k+1$ . Let  $\Phi_1 := \Phi_{D_1}(s, X)$ ,  $\Phi_2 := \Phi_{D_2}(s, X)$ . Then the sums defining  $\Phi_1$  and  $\Phi_2$  differ only in the  $m$ -th and  $m+1$ -th term, so we get

$$\Phi_1 - \Phi_2 =$$

$$\varphi(YabZ[c, d]T) + \varphi(Y[a, b]ZdcT) - \varphi(Y[a, b]ZcdT) - \varphi(YbaZ[c, d]T),$$

which equals zero by the induction assumption.

For the third relation, suppose that we have two decompositions  $D_1, D_2$  of  $s$  given by  $s = ps_j s_{j+1} s_j q$  and  $s = ps_{j+1} s_j s_{j+1} q$ , where  $q$  is a product of  $k$  transpositions of neighbors. Let  $q(X) = YabcZ$  where  $a, b, c \in \mathfrak{g}$  stand in positions  $j, j+1, j+2$ . Let  $\Phi_1 := \Phi_{D_1}(s, X)$ ,  $\Phi_2 := \Phi_{D_2}(s, X)$ . Then the sums defining  $\Phi_1$  and  $\Phi_2$  differ only in the  $k$ -th,  $k+1$ -th, and  $k+2$ -th terms, so we get

$$\Phi_1 - \Phi_2 =$$

$$\begin{aligned} & (\varphi(Y[a, b]cZ) + \varphi(Yb[a, c]Z) + \varphi(Y[b, c]aZ)) - \\ & (\varphi(Ya[b, c]Z) + \varphi(Y[a, c]bZ) + \varphi(Yc[a, b]Z)). \end{aligned}$$

So the Jacobi identity

$$[[b, c], a] + [b, [a, c]] + [[a, b], c] = 0$$

combined with property (ii) in degree  $n-1$  implies that  $\Phi_1 - \Phi_2 = 0$ , i.e.,  $\Phi_1 = \Phi_2$ , as claimed. Thus we will denote  $\Phi_D(s, X)$  just by  $\Phi(s, X)$ .

It remains to show that  $\Phi(s, X)$  does not depend on the choice of  $s$  and only depends on  $s(X)$ . Let  $X = x_{i_1} \dots x_{i_n}$ ; then  $s(X) = s'(X)$  if and only if  $s = s't$ , where  $t$  is the product of transpositions  $s_k$  for which  $i_k = i_{k+1}$ . Thus, it suffices to show that  $\Phi(s, X) = \Phi(ss_k, X)$  for such  $k$ . But this follows from the fact that  $[x, x] = 0$ .

Now, it follows from the construction of  $\varphi$  that for any monomial  $X$  of degree  $n$  (not necessarily ordered),  $\varphi(s_j(X)) = \varphi(X) + \varphi([, ]_j(X))$ .

Thus  $\varphi$  satisfies property (ii) in degree  $n$ . This concludes the proof of Lemma 13.11 and hence Theorem 13.1.  $\square$

MIT OpenCourseWare  
<https://ocw.mit.edu>

18.745 Lie Groups and Lie Algebras I  
Fall 2020

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.