

Complex Numbers

- Complex numbers have both real and imaginary components. A complex number \underline{r} may be expressed in Cartesian or Polar forms:

$$\begin{aligned}\underline{r} &= a + jb \text{ (cartesian)} \\ &= |r|e^{j\phi} \text{ (polar)}\end{aligned}$$

The following relationships convert from cartesian to polar forms:

$$\begin{aligned}\text{Magnitude } |r| &= \sqrt{a^2 + b^2} \\ \text{Angle } \phi &= \begin{cases} \tan^{-1} \frac{b}{a} & a > 0 \\ \tan^{-1} \frac{b}{a} \pm \pi & a < 0 \end{cases}\end{aligned}$$

- Complex numbers can be plotted on the complex plane in either Cartesian or Polar forms Fig.1.




Figure 1: Complex plane plots: Cartesian and Polar forms

Euler's Identity

Euler's Identity states that

$$e^{j\phi} = \cos \phi + j \sin \phi$$

This can be shown by taking the series expansion of \sin , \cos , and e .

$$\begin{aligned}\sin \phi &= \phi - \frac{\phi^3}{3!} + \frac{\phi^5}{5!} - \frac{\phi^7}{7!} + \dots \\ \cos \phi &= 1 - \frac{\phi^2}{2!} + \frac{\phi^4}{4!} - \frac{\phi^6}{6!} + \dots \\ e^{j\phi} &= 1 + j\phi - \frac{\phi^2}{2!} - j\frac{\phi^3}{3!} + \frac{\phi^4}{4!} + j\frac{\phi^5}{5!} + \dots\end{aligned}$$

Combining

$$\begin{aligned}\cos \phi + j \sin \phi &= 1 + j\phi - \frac{(\phi)^2}{2!} - j\frac{\phi^3}{3!} + \frac{\phi^4}{4!} + j\frac{\phi^5}{5!} + \dots \\ &= e^{j\phi}\end{aligned}$$

Complex Exponentials

- Consider the case where ϕ becomes a function of time increasing at a constant rate ω

$$\phi(t) = \omega t.$$

then $\underline{r}(t)$ becomes

$$\underline{r}(t) = e^{j\omega t}$$



Plotting $\underline{r}(t)$ on the complex plane traces out a circle with a constant radius = 1 (Fig. 2). Plotting the real and imaginary components of $\underline{r}(t)$ vs time (Fig. 3), we see that the real component is $Re\{\underline{r}(t)\} = \cos \omega t$ while the imaginary component is $Im\{\underline{r}(t)\} = \sin \omega t$.

- Consider the variable $\underline{r}(t)$ which is defined as follows:

$$\underline{r}(t) = e^{\underline{s}t}$$

where \underline{s} is a complex number

$$\underline{s} = \sigma + j\omega$$

Figure 2: Complex plane plots: $\underline{r}(t) = e^{j\omega t}$ Figure 3: Real and imaginary components of $\underline{r}(t)$ vs time

- What path does $\underline{r}(t)$ trace out in the complex plane? Consider

$$\underline{r}(t) = e^{st} = e^{(\sigma + j\omega)t} = e^{\sigma t} \cdot e^{j\omega t}$$

One can look at this as a time varying magnitude ($e^{\sigma t}$) multiplying a point rotating on the unit circle at frequency ω via the function $e^{j\omega t}$. Plotting just the magnitude of $e^{j\omega t}$ vs time shows that there are three distinct regions (Fig. 4):

1. $\sigma > 0$ where the magnitude grows without bounds. This condition is unstable.
2. $\sigma = 0$ where the magnitude remains constant. This condition is

called marginally stable since the magnitude does not grow without bound but does not converge to zero.

3. $\sigma < 0$ where the magnitude converges to zero. This condition is termed stable since the system response goes to zero as $t \rightarrow \infty$.




Figure 4: Magnitude $r(t)$ for various σ .




Effect of Pole Position

The stability of a system is determined by the location of the system poles. If a pole is located in the *2nd* or *3rd* quadrant (which quadrant determines the direction of rotation in the polar plot), the pole is said to be stable.

Figure 5 shows the pole position in the complex plane, the trajectory of $\underline{r}(t)$ in the complex plane, and the real component of the time response for a stable pole.

If the pole is located directly on the imaginary axis, the pole is said to be marginally stable. Figure 6 shows the pole position in the complex plane, the trajectory of $\underline{r}(t)$ in the complex plane, and the real component of the time response for a marginally stable pole.

Lastly, if a pole is located in either the *1st* or *4th* quadrant, the pole is said to be unstable. Figure 7 shows the pole position in the complex plane, the trajectory of $\underline{r}(t)$ in the complex plane, and the real component of the time response for an unstable pole.

Figure 5: Pole position, $\underline{r}(t)$, and real time response for stable pole.Figure 6: Pole position, $\underline{r}(t)$, and real time response for marginally stable pole.Figure 7: Pole position, $\underline{r}(t)$, and real time response for unstable pole.