


WAVEMAKER THEORY




A PADDLE WITH DRAFT D IS UNDERGOING SMALL AMPLITUDE HORIZONTAL OSCILLATIONS WITH DISPLACEMENT

$$\xi(t) = \text{Re}\{\Xi e^{i\omega t}\}$$


WHERE Ξ IS ASSUMED KNOWN AND REAL.

THIS EXCITATION CREATES PLANE PROGRESSIVE WAVES WITH AMPLITUDE A DOWN THE TANK. THE PRINCIPAL OBJECTIVE OF WAVEMAKER THEORY IS TO DETERMINE A AS A FUNCTION OF ω , Ξ AND H .

OTHER TYPES OF WAVEMAKER MODES, LIKE



OR



OR



MAY BE TREATED SIMILARLY.

IN GENERAL, THE WAVEMAKER DISPLACEMENT AT $x=0$ MAY BE WRITTEN IN THE FORM

$$\zeta(t) = \text{Re} \{ \mathbb{H}(z) e^{i\omega t} \}$$

WHERE $\mathbb{H}(z)$ IS A KNOWN FUNCTION OF z . —

LET THE TOTAL VELOCITY POTENTIAL BE:

$$\phi = \text{Re} \{ \varphi e^{i\omega t} \}$$

WHERE

$$\varphi = \varphi_w + \psi$$

THE FIRST TERM IS A VELOCITY POTENTIAL THAT REPRESENTS A PLANE PROGRESSIVE REGULAR WAVE DOWN THE TANK WITH AMPLITUDE A , YET UNKNOWN. THUS:

$$\varphi_w = \frac{igA}{\omega} \frac{\cosh k(z+H)}{\cosh kH} e^{-ikx+i\omega t}$$

WITH :

$$\omega^2 = gk \tanh kH. —$$

THE SECOND COMPONENT POTENTIAL ψ IS BY DEFINITION A DECAYING DISTURBANCE AS $x \rightarrow \infty$ AND OTHERWISE SATISFIES THE FOLLOWING BOUNDARY VALUE PROBLEM:

$$\left\{ \begin{array}{l} \nabla^2 \psi = \psi_{xx} + \psi_{zz} = 0, \quad -H < z < 0 \\ \psi_z - \frac{\omega^2}{g} \psi = 0, \quad z = 0 \\ \psi_z = 0, \quad z = -H \\ \psi \rightarrow 0, \quad x \rightarrow \infty \end{array} \right.$$

THE CONDITION ON THE WAVEMAKER ($x=0$) IS YET TO BE ENFORCED.

NOTE THAT UNLIKE ϕ_w , ψ IS NOT REPRESENTING A PROPAGATING WAVE DOWN THE TANK SO IT IS CALLED A NON-WAVELIKE MODE. SUCH MODES DO EXIST AS WILL BE SHOWN BELOW. ON THE WAVEMAKER ($x=0$) THE HORIZONTAL VELOCITY DUE TO ϕ_w AND THAT DUE TO ψ MUST SUM TO THE FORCING VELOCITY DUE TO $\xi(t)$. —

NOTING THAT $\varphi_w \sim e^{-ikx} \cosh k(z+H)$

WE WILL TRY $\psi \sim e^{-\lambda x} \cos \lambda(z+H)$

ITS CONJUGATE WHICH SATISFIES THE
CONDITION OF VANISHING VALUE AS $x \rightarrow \infty$
FOR $\lambda > 0$. —

LAPLACE: $\psi_{xx} + \psi_{zz} = 0$, VERIFY FOR
ALL λ

FS CONDITION: $\psi_z - \frac{\omega^2}{g} \psi = 0 \Rightarrow$
 $-\lambda \sin \lambda H - \frac{\omega^2}{g} \cos \lambda H = 0$
 $\Rightarrow \lambda \tan \lambda H = -\nu \equiv \frac{\omega^2}{g}$

SEAFLOOR } $\psi_z = 0$, $z = -H$, (VERIFY)
CONDITION }

SO FOR THE NON-WAVELIKE MODES ψ , λ
MUST SATISFY THE "DISPERSION" RELATION


$$\lambda \tan \lambda H = -\nu = -\frac{\omega^2}{g} < 0$$

FOR POSITIVE VALUES OF λ SO THAT

$e^{-\lambda x} \rightarrow 0$, $x \rightarrow +\infty$. —

VALUES OF ω_i SATISFYING THE DISPERSION RELATION FOLLOW FROM THE SOLUTION OF THE NON-DIMENSIONAL NONLINEAR EQUATION

$$\tan w = -\frac{\nu}{w}, \quad w = \omega H$$



SOLUTIONS ω_i , $i=1, 2, \dots$ EXIST AS SHOWN ABOVE WITH $\omega_i \sim i\pi$ FOR LARGE i . THESE VALUES ARE KNOWN AS THE EIGENVALUES OR EIGEN-WAVENUMBERS OF THE NON-WAVELIKE MODES. THE EIGEN-WAVENUMBER OF THE WAVELIKE SOLUTION k IS GIVEN BY THE DISPERSION RELATION:

$$\frac{\omega^2 H}{g} = kH \tan kH.$$

VERIFY THAT BY SETTING $K = i\lambda$;
 THE DISPERSION RELATION OF THE
 NON-WAVELIKE MODES FOLLOWS. IN
 SUMMARY THE PURELY IMAGINARY ROOTS
 OF THE SURFACE WAVE DISPERSION
 RELATION AND ITS SINGLE REAL POSITIVE
 ROOT ENTER THE SOLUTION OF THE
 WAVEMAKER PROBLEM.

DEFINE THE FOLLOWING ORTHOGONAL
 EIGENMODES IN THE VERTICAL DIRECTION z :

$$f_0(z) = \frac{\sqrt{2} \cosh K(z+H)}{(H + \frac{1}{2} \sinh^2 KH)^{1/2}}$$

$$f_n(z) = \frac{\sqrt{2} \cos \lambda_n (z+H)}{(H - \frac{1}{2} \sin^2 \lambda_n H)}, \quad n=1, 2, \dots$$

SELECTED TO SATISFY :

$$\left\{ \begin{array}{l} \int_{-H}^0 f_0^2(z) dz = \int_{-H}^0 f_n^2(z) dz = 1 \\ \int_{-H}^0 f_m(z) f_n(z) dz = 0, \quad m \neq n \end{array} \right.$$

SO THE WAVEMAKER VELOCITY POTENTIALS ϕ_w AND ψ CAN BE EXPRESSED SIMPLY IN TERMS OF THEIR RESPECTIVE EIGENMODES:

$$\phi_w = a_0 f_0(z) e^{-ikx}$$

$$\psi = \sum_{n=1}^{\infty} a_n f_n(z) e^{-\lambda_n x}$$

AND:

$$\phi = \text{Re} \left\{ (\phi_w + \psi) e^{i\omega t} \right\}$$

ON $x=0$:

$$\phi_x = \text{Re} \left\{ (\phi_w + \psi_x) \Big|_{x=0} e^{i\omega t} \right\}$$

$$= \frac{d\xi}{dt} = \text{Re} \left\{ \mathbb{H}(z) i\omega e^{i\omega t} \right\}$$

OR

$$\frac{\partial}{\partial x} (\phi_w + \psi) \Big|_{x=0} = \mathbb{H}(z) i\omega$$

↑
known

$$\frac{\partial \phi_w}{\partial x} \Big|_{x=0} = a_0 (-ik) f_0(z)$$

$$\frac{\partial \psi}{\partial x} \Big|_{x=0} = \sum_{n=1}^{\infty} a_n (-\lambda_n) f_n(z)$$

IT FOLLOWS THAT:

$$-ik a_0 f_0(z) + \sum_{n=1}^{\infty} a_n (-\lambda_n) f_n(z) = i\omega \mathbb{F}(z)$$

- ONE OF THE PRIMARY OBJECTIVES OF WAVEMAKER THEORY IS TO DETERMINE a_0 (OR THE FAR-FIELD WAVE AMPLITUDE A) IN TERMS OF $\mathbb{F}(z)$. MULTIPLYING BOTH SIDES BY $f_0(z)$, INTEGRATING FROM $-H \rightarrow 0$ AND USING ORTHOGONALITY WE OBTAIN:

$$-ik a_0 = i\omega \int_{-H}^0 dz f_0(z) \mathbb{F}(z)$$

$$\Rightarrow a_0 = -\frac{\omega}{k} \int_{-H}^0 dz f_0(z) \mathbb{F}(z)$$

THE FAR-FIELD WAVE COMPONENT REPRESENTING PROPAGATING WAVES IS GIVEN BY:

$$\varphi_w = a_0 \frac{\sqrt{2} \cosh k(z+H)}{(H + \frac{1}{2} \sinh^2 kH)^{1/2}} e^{-ikx}$$

$$\equiv \frac{igA}{\omega} \frac{\cosh k(z+H)}{\cosh kH} e^{-ikx}$$


PLUGGING IN a_0 AND SOLVING FOR A
 WE OBTAIN THE COMPLEX AMPLITUDE OF
 THE PROPAGATING WAVE AT INFINITY,
 NAMELY MODULUS AND PHASE, IN TERMS
 OF THE WAVE MAKER DISPLACEMENT
 $\mathbb{H}(z)$ AND THE OTHER FLOW PARAMETERS.

EXERCISES

- TRY $\mathbb{H}(z) = A \begin{cases} 1, & -D < z < 0 \\ 0, & -H < z < -D \end{cases}$

PADDLE-TYPE WAVEMAKER. DETERMINE
 THE AMPLITUDE A_w AND PHASE OF
 THE FAR-FIELD WAVE-TRAIN.

- REPEAT ABOVE EXERCISE FOR A HINGE
 TYPE WAVEMAKER?




- FOR WHAT TYPE OF $\mathbb{H}(z)$ ARE THE NON-WAVELIKE MODES $\psi \equiv 0$? IT IS EASY TO VERIFY BY VIRTUE OF ORTHOGONALITY THAT:

$$\mathbb{H}(z) \approx f_0(z)$$

UNFORTUNATELY THIS IS NOT A "PRACTICAL" DISPLACEMENT SINCE $f_0(z, k)$ DEPENDS ON k , THUS ON ω . SO ONE WOULD NEED TO BUILD A FLEXIBLE PADDLE!

- WHAT IS THE WAVE AMPLITUDE AT INFINITY GENERATED BY A POINT SOURCE LOCATED AT $z = -D$?



MORE DETAILS ON THE ABOVE THEORY AND EXTENSIONS TO THE NONLINEAR CASE MAY BE FOUND IN W&L AND MEI. —