


20.330J Fields, Forces and Flows in Biological Systems
 Prof. Scott Manalis
Review: Vector Calculus

Vector Product

$$\vec{v} \cdot \hat{n} = v_x n_x + v_y n_y + v_z n_z$$

$$= |\vec{v}| |\hat{n}| \cos(\alpha)$$



Gradient (on a scalar function)

$$\vec{\nabla} = \hat{i}_x \frac{\partial}{\partial x} + \hat{i}_y \frac{\partial}{\partial y} + \hat{i}_z \frac{\partial}{\partial z}$$

$$\vec{\nabla} p = \hat{i}_x \frac{\partial p}{\partial x} + \hat{i}_y \frac{\partial p}{\partial y} + \hat{i}_z \frac{\partial p}{\partial z}$$

Divergence (operated on vector)


$$\vec{\nabla} \cdot \vec{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z} \quad \Rightarrow \text{scalar}$$

Curl (operated on vector)

$$\vec{\nabla} \times \vec{v} = \begin{vmatrix} \hat{i}_x & \hat{i}_y & \hat{i}_z \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ v_x & v_y & v_z \end{vmatrix} \quad \Rightarrow \text{vector}$$

In 1D integration...

$$f(x_2) - f(x_1) = \int_{x_1}^{x_2} \frac{df}{dx} dx$$



...similarly, we have two different integral theorems for vector calculus.

(1) Gauss' theorem (Divergence theorem)

For any vector field \hat{v} ,


$$\oint_S \hat{v} \cdot \hat{n} da = \int_V (\nabla \cdot \hat{v}) dv$$

velocity \times *area*


"total outgoing volume flow rate"

"volume expansion"

surface S



Proof: consider infinitesimal cube.



From surfaces ① and ②:

$$\oint_S (\hat{v} \cdot \hat{n}) da \rightarrow (V_x|_{x+\Delta x} - V_x|_x) \Delta y \Delta z$$

Similarly, from other surfaces,


$$\begin{aligned} \oint_S (\hat{v} \cdot \hat{n}) da &= (V_x|_{x+\Delta x} - V_x|_x) \Delta y \Delta z \\ &\quad + (V_y|_{y+\Delta y} - V_y|_y) \Delta x \Delta z \\ &\quad + (V_z|_{z+\Delta z} - V_z|_z) \Delta x \Delta y \end{aligned}$$

Divide each terms with Δx , Δy , Δz respectively,

$$\begin{aligned} &= \left[\frac{\partial V_x}{\partial x} + \frac{\partial V_y}{\partial y} + \frac{\partial V_z}{\partial z} \right] \Delta x \Delta y \Delta z \\ &= \oint_V (\nabla \cdot \vec{V}) dV \end{aligned}$$

Meaning of “ $\nabla \cdot \vec{V}$ ”


- volume expansion
- net outgoing flux
- for incompressible flow, $\nabla \cdot \vec{V} = 0$ (no fluid source/sink)



(2) Stokes' theorem (curl theorem)

For a given vector field \hat{v} ,

$$\oint_C \vec{V} \cdot d\vec{s} = \int_S (\nabla \times \vec{V}) \cdot \hat{n} da$$




Proof: think about the rectangle in the xy plane.

$$\begin{aligned} \oint_C \vec{V} \cdot d\vec{s} &= (V_x|_y - V_x|_{y+\Delta y})\Delta x + (V_y|_{x+\Delta x} - V_y|_x)\Delta y \\ &= \left(-\frac{V_x|_{y+\Delta y} - V_x|_y}{\Delta y} + \frac{V_y|_{x+\Delta x} - V_y|_x}{\Delta x} \right) \Delta x \Delta y \\ &= \left(\frac{\partial V_y}{\partial x} - \frac{\partial V_x}{\partial y} \right) \Delta x \Delta y = (\nabla \times \vec{V})_z \Delta x \Delta y \end{aligned}$$


Similar for curves in other planes...

Meaning of “ $\nabla \times \vec{V}$ ”



- Represents “circulation” of the flow.



$\nabla \times \vec{V} = 0$
Laminar flow



$\nabla \times \vec{V} \neq 0$
Turbulent flow



References

- H&M website: Chapter 2
- Appendix of TY & K