

## The following graph of $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$ describes a reaction in which:



- 1.  $\Delta H^{\circ} > 0$  and  $\Delta S^{\circ} > 0$
- 2.  $\Delta H^{\circ} < 0$  and  $\Delta S^{\circ} < 0$
- 3.  $\Delta H^{\circ} < 0$  and  $\Delta S^{\circ} > 0$
- 4.  $\Delta H^{\circ} > 0$  and  $\Delta S^{\circ} < 0$

1

## The following graph of $\Delta G^{\circ} = \Delta H^{\circ} - T\Delta S^{\circ}$ describes a reaction in which:



1. 
$$\Delta H^{\circ} > 0$$
 and  $\Delta S^{\circ} > 0$ 

73% 
$$\Delta H^{\circ} < 0$$
 and  $\Delta S^{\circ} < 0$ 

3. 
$$\Delta H^{\circ} < 0$$
 and  $\Delta S^{\circ} > 0$ 

8%

4. 
$$\Delta H^{\circ} > 0$$
 and  $\Delta S^{\circ} < 0$ 

$$N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$$

$$K = 1.9 \times 10^{-4} \text{ at } 400^{\circ}\text{C}$$
  
 $PN_2 = 5.5 \text{ bar}, PH_2 = 2.2 \text{ bar}, PNH_3 = 1.1 \text{ bar at } 400^{\circ}\text{C}$ 

#### Which direction will the reaction go?

- 1. toward products, since Q < K
- 2. toward products, since Q > K
- 3. toward reactants, since Q > K
- 4. toward reactants, since Q < K

$$N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$$

$$K = 1.9 \times 10^{-4} \text{ at } 400^{\circ}\text{C}$$
  
 $PN_2 = 5.5 \text{ bar, } PH_2 = 2.2 \text{ bar, } PNH_3 = 1.1 \text{ bar at } 400^{\circ}\text{C}$ 

#### Which direction will the reaction go?

- 1. toward products, since Q < K
- 2. toward products, since Q > K
- 79%  $\odot$ 3. toward reactants, since Q > K
- 4. toward reactants, since Q < K

# Which is the correct expression for K for the following reaction:

$$2NO_2$$
 (g)  $\rightarrow N_2O_4$ (g)

- 1.  $(PNO_2)/(PN_2O_4)$
- 2.  $(PNO_2)^2/(PN_2O_4)$
- 3.  $(PN_2O_4)/(PNO_2)$
- 4.  $(PN_2O_4)/(PNO_2)^2$

# Which is the correct expression for K for the following reaction:

$$2NO_2$$
 (g)  $\rightarrow N_2O_4$ (g)

- $1. (PNO_2)/(PN_2O_4)$
- 8% 2.  $(PNO_2)^2/(PN_2O_4)$
- 3% 3.  $(PN_2O_4)/(PNO_2)$
- 88%  $\sqrt{4}$ .  $(PN_2O_4)/(PNO_2)^2$

### $\Delta G^{\circ} = -RT \ln K$ or $K = \exp [-\Delta G^{\circ} / RT]$ K is large if $\Delta G^{\circ}$ is...

- 1. negative and small
- 2. negative and large
- 3. positive and small
- 4. positive and large

### $\Delta \mathbf{G}^{\circ} = -\mathbf{R}\mathbf{T} \ln \mathbf{K}$ or $\mathbf{K} = \exp \left[-\Delta \mathbf{G}^{\circ} / \mathbf{R}\mathbf{T}\right]$ K is large if $\Delta \mathbf{G}^{\circ}$ is...

1. negative and small

82% 2. negative and large

5% 3. positive and small

4. positive and large


#### **Removing Product**

### If you remove product, what happens?

- 1. Q<K. The reaction shifts to the right toward product.
- 2. Q>K. The reaction shifts to the right toward product.
- 3. Q<K. The reaction shifts to the left toward reactants.
- 4. Q>K. The reaction shifts to the left toward reactants.

#### **Removing Product**

If you remove product, what happens?



1. Q<K. The reaction shifts to the right toward product.



2. Q>K. The reaction shifts to the right toward product.



3. Q<K. The reaction shifts to the left toward reactants.



4. Q>K. The reaction shifts to the left toward reactants.

MIT OpenCourseWare http://ocw.mit.edu

5.111 Principles of Chemical Science Fall 2014

For information about citing these materials or our Terms of Use, visit: http://ocw.mit.edu/terms.