

Problem Set 4
Due October 6, 2010

1. Random variables X and Y have the joint PMF

$$p_{X,Y}(x, y) = \begin{cases} c(x^2 + y^2), & \text{if } x \in \{1, 2, 4\} \text{ and } y \in \{1, 3\}, \\ 0, & \text{otherwise.} \end{cases}$$

- (a) What is the value of the constant c ?
 - (b) What is $\mathbf{P}(Y < X)$?
 - (c) What is $\mathbf{P}(Y > X)$?
 - (d) What is $\mathbf{P}(Y = X)$?
 - (e) What is $\mathbf{P}(Y = 3)$?
 - (f) Find the marginal PMFs $p_X(x)$ and $p_Y(y)$.
 - (g) Find the expectations $\mathbf{E}[X]$, $\mathbf{E}[Y]$ and $\mathbf{E}[XY]$.
 - (h) Find the variances $\text{var}(X)$, $\text{var}(Y)$ and $\text{var}(X + Y)$.
 - (i) Let A denote the event $X \geq Y$. Find $\mathbf{E}[X | A]$ and $\text{var}(X | A)$.
2. The newest invention of the 6.041/6.431 staff is a three-sided die with faces numbered 1, 2, and 3. The PMF for the result of any one roll of this die is

$$p_X(x) = \begin{cases} 1/2, & \text{if } x = 1, \\ 1/4, & \text{if } x = 2, \\ 1/4, & \text{if } x = 3, \\ 0, & \text{otherwise.} \end{cases}$$

Consider a sequence of six independent rolls of this die, and let X_i be the random variable corresponding to the i th roll.

- (a) What is the probability that exactly three of the rolls have result equal to 3?
 - (b) What is the probability that the first roll is 1, given that exactly two of the six rolls have result of 1?
 - (c) We are told that exactly three of the rolls resulted in 1 and exactly three resulted in 2. Given this information, what is the probability that the sequence of rolls is 121212?
 - (d) Conditioned on the event that at least one roll resulted in 3, find the conditional PMF of the number of 3's.
3. Suppose that X and Y are independent, identically distributed, geometric random variables with parameter p . Show that

$$\mathbf{P}(X = i | X + Y = n) = \frac{1}{n-1}, \quad \text{for } i = 1, 2, \dots, n-1.$$

MASSACHUSETTS INSTITUTE OF TECHNOLOGY
Department of Electrical Engineering & Computer Science
6.041/6.431: Probabilistic Systems Analysis
(Fall 2010)

4. Consider 10 independent tosses of a biased coin with a probability of heads of p .
- (a) Let A be the event that there are 6 heads in the first 8 tosses. Let B be the event that the 9th toss results in heads. Show that events A and B are independent.
 - (b) Find the probability that there are 3 heads in the first 4 tosses and 2 heads in the last 3 tosses.
 - (c) Given that there were 4 heads in the first 7 tosses, find the probability that the 2nd head occurred during the 4th trial.
 - (d) Find the probability that there are 5 heads in the first 8 tosses and 3 heads in the last 5 tosses.
5. Consider a sequence of independent tosses of a biased coin at times $t = 0, 1, 2, \dots$. On each toss, the probability of a 'head' is p , and the probability of a 'tail' is $1 - p$. A reward of one unit is given each time that a 'tail' follows immediately after a 'head.' Let R be the total reward paid in times $1, 2, \dots, n$. Find $\mathbf{E}[R]$ and $\text{var}(R)$.

G1[†]. A simple example of a random variable is the *indicator* of an event A , which is denoted by I_A :

$$I_A(\omega) = \begin{cases} 1, & \text{if } \omega \in A \\ 0, & \text{otherwise.} \end{cases}$$

- (a) Prove that two events A and B are independent if and only if the associated indicator random variables, I_A and I_B are independent.
- (b) Show that if $X = I_A$, then $\mathbf{E}[X] = \mathbf{P}(A)$.

[†]Required for 6.431; optional for 6.041

MIT OpenCourseWare
<http://ocw.mit.edu>

6.041SC Probabilistic Systems Analysis and Applied Probability
Fall 2013

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.