


An Example of Clustering Expression Data

6.874 / 7.90


Computational Functional Genomics

Spring 2005


Results – hand generated data




Hierarchical clustering




Input




Optimal ordering



Hierarchical clustering



Input



Optimal ordering

Problem definition

Denote by Φ the space of the possible linear orderings consistent with the tree.


Denote by $v_1 \dots v_n$ the tree leaves.

Our goal is to find an ordering that maximizes the similarity of adjacent elements:

$$\max_{\phi \in \Phi} \sum_{i=1}^{n-1} S(v_i^\phi, v_{i+1}^\phi)$$

where S is the similarity matrix

59 experiments,
combining cdc15,
cdc28 and α factor
arrest



Cluster


G2/M

G1

M/G1

S/G2

S



Optimal

G2/M

S/G2

S

M/G1

G1

M/G1

Discovering new types of cancer

Alizadeh, A., Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling, Nature 403, pp. 503-511 (February 3, 2000)

(Does not use optimal ordering)

Goal

- Discover cause for different disease courses for diffuse large B-cell lymphoma (DLBCL)
 - 40% of patients respond to therapy
 - 60% succumb to disease
- Provide diagnostic / prognostic tool
- DLBCL is most common subtype of non-Hodgkin's lymphoma

Questions

- Can we create a molecular portrait of distinct types of B-cell malignancy?
- Can we identify types of malignancy not yet recognized?
- Can we relate malignancy to normal stages in B-cell development and physiology?

Lymphochip

- 17,856 cDNA clones
 - 12,069 from germinal B-cell library
 - 2,338 from DLBCL, follicular lymphoma (FL), mantle cell lymphoma, and chronic lymphocytic leukaemia (CLL)
 - 3,186 genes important to lymphocyte and/or cancer biology
 - B- and T-lymphocyte genes that respond to mitogens or cytokines

Data sources

- Rearranged immunoglobulin genes in DLBCL are characteristic of germinal center of secondary lymphoid organs
- 96 normal and malignant lymphocyte samples

Figures removed for copyright reasons.

See Alizadeh, A., et al. "Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling." *Nature* 403 (February 3, 2000): 503-511.

Summary

- DLBCL groups are still diverse – some members of GC B-like DLBCL group die
 - 5 in first 2 years
- May be able to find informative features for more groups
- If can find constitutive genes in cancers, target upstream regulators