

8.324 Relativistic Quantum Field Theory II

MIT OpenCourseWare Lecture Notes

Hong Liu, Fall 2010

Lecture 18

4.2: CANCELLATION OF DIVERGENCES

4.2.1: General Structure

Consider a generic divergent diagram M of degree D ; that is,

$$M \sim \int^\Lambda ds s^{D-1} \quad (1)$$

if all loop momenta are taken proportional to s . Generically, internal propagators have the form

$$\frac{1}{(as + p)^\alpha + \dots} \sim \frac{1}{s^\alpha} \quad \alpha = \begin{cases} 2 & \text{scalar or vector,} \\ 1 & \text{fermion.} \end{cases} \quad (2)$$

for large s , where a is a numerical constant and p is a combination of the external momenta. Differentiating M , with respect to p , n times gives a term proportional to

$$\frac{1}{(as + p)^{\alpha+n}} \sim \frac{1}{s^{\alpha+n}}, \quad (3)$$

and so $D + 1$ derivatives with respect to the external momenta will make M finite. This means that

$$M(p) = M_0 + M_1 p + \dots + M_D p^D + \text{finite terms}, \quad (4)$$

where the argument p of the function represents the collection of external momenta, we have suppressed the index structure, and M_0, M_1, \dots, M_D are potentially divergent constants. Suppose that M has E_f external lines of the field ϕ_f . Then divergences of $M(p)$ can be cancelled by counterterms of the form

$$\sum_{j=0}^D A_j (\partial)^j \prod_f \phi_f^{E_f} \quad (5)$$

where the A_j are divergent coefficients in order to cancel divergences in M_j . The index structure in $A_j \partial^j$ should match the suppressed index structure of $M(p)$.

4.2.2: Renormalization of Quantum Electrodynamics

In the case of quantum electrodynamics,

$$D = 4 - E_\gamma - \frac{3}{2} E_e, \quad (6)$$

and hence, ultraviolet-divergent diagrams are given by

$$E_\gamma + \frac{3}{2} E_e \leq 4. \quad (7)$$

We now enumerate these classes of diagrams:

1. $E_\gamma = E_e = 0, D = 4$: These are the vacuum diagrams, and give a divergent constant Λ_0 . The divergence can be removed by including a constant Λ_{ct} counterterm in the Lagrangian, such that

$$\Lambda_{phys} = \Lambda_0 - \Lambda_{ct}. \quad (8)$$

Λ_{phys} does not have any physical significance, unless we are dealing with a theory of gravity. Λ_{phys} is measured to be very small, and it acts as a cosmological constant.

where A is a logarithmically divergent coefficient. This can be cancelled by the counterterm three-point vertex,

$$\begin{array}{c} \diagup \\ \text{---} \times \text{---} \\ \diagdown \end{array} = -Z_1 e A_\mu \bar{\psi} \gamma^\mu \psi, \tag{19}$$

where the Ward identity imposes $Z_1 = Z_2$.

In summary, the counter terms we considered before are enough to cancel all of the superficial divergences to all orders in perturbation theory. We still need to consider diagrams with divergent sub-diagrams. Generically, these can be easily cancelled since divergent sub-graphs must also belong to those considered above. For example, the pairings

$$\begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array}, \quad \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array} \tag{20}$$

are both finite. We now need to consider the case of overlapping divergences. That is, when two divergent subgraphs share an internal line. These divergences are more difficult to deal with. For example, consider the diagram

$$\begin{array}{c} p+q \qquad p'+q \\ \text{---}^\mu \qquad \text{---}^\nu \\ q \qquad q \\ p \qquad p' \end{array} \tag{21}$$

In the case that p is taken to be large, the pairing

$$\begin{array}{c} \text{---}^\mu \qquad \text{---}^\nu \\ q \qquad q \\ p' \end{array} + \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array} \tag{22}$$

gives a vertex correction at the vertex μ , leaving a residual self-energy divergence for the case $p' \rightarrow \infty$. Similarly, if p' is taken to be large, the pairing

$$\begin{array}{c} \text{---}^\mu \qquad \text{---}^\nu \\ q \qquad q \\ p \end{array} + \begin{array}{c} \text{---} \times \text{---} \\ \text{---} \end{array} \tag{23}$$

gives a vertex correction at the vertex ν , leaving a residual self-energy divergence for the case $p \rightarrow \infty$. The two procedures are not independent, and it is finally necessary to add a correction of the form

$$\text{---} \times \text{---} \tag{24}$$

at the order $O(e^4)$ to cancel the divergences when both p and p' are large. Another example is given by the combination of diagrams

$$\text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} + \text{Diagram 4} \quad (25)$$

It can be proved to all orders, the content of the Bogoliubov-Parasiuk-Hepp-Zimmermann theorem, that overlapping divergences can be canceled by counterterms corresponding to superficial divergences.

MIT OpenCourseWare
<http://ocw.mit.edu>

8.324 Relativistic Quantum Field Theory II
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.