

**Solutions Day 7, F 3/1/2024**  
 Topic 4: Complex numbers (day 2 of 2)  
 Jeremy Orloff

**Problem 1.** Let  $z_1 = 2 + 5i$ ,  $z_2 = 1 + 3i$

(a) Compute  $z_1 + z_2$ ,  $z_1 \cdot z_2$ ,  $z_1 \cdot \bar{z}_1$ ,  $|z_1|$ ,  $\text{Arg}(z_1)$ .

**Solution:**  $z_1 + z_2 = 3 + 8i$ .

$$z_1 \cdot z_2 = (2 + 5i)(1 + 3i) = 2 - 15 + 6i + 5i = -13 + 11i.$$

$$z_1 \cdot \bar{z}_1 = (2 + 5i)(2 - 5i) = 4 + 25 = 29.$$

$$|z_1| = \sqrt{2^2 + 5^2} = \sqrt{29}.$$

$$\text{Arg}(z_1) = \tan^{-1}(5/2) \text{ in Q1}.$$

(b) Find  $\text{Re}(z_1)$ ,  $\text{Im}(z_1)$ .

**Solution:**  $\text{Re}(z_1) = 2$ ,  $\text{Im}(z_1) = 5$ . (Imaginary part is a real number!)

(c) Let  $z = x + iy$ . Compute  $z \cdot \bar{z}$ .


$$\text{Solution: } z \cdot \bar{z} = (x + iy)(x - iy) = x^2 + y^2 = |z|^2.$$

**Problem 2.** Find  $r$  and  $\theta = \text{Arg}(z)$  for  $z = i$ ,  $z = 1 - i$ ,  $z = 5(1 - i)$ .

**Solution:**  $z = i$ :  $|i| = 1$ ,  $\theta = \text{Arg}(i) = \pi/2 + 2n\pi$ ,  $n$  an integer.

$z = 1 - i$ :  $|1 - i| = \sqrt{2}$ ,  $\theta = \text{Arg}(1 - i) = -\pi/4 + 2n\pi$ ,  $n$  an integer.


$z = 5(1 - i)$ :  $|5(1 - i)| = 5\sqrt{2}$ ,  $\theta = \text{Arg}(5(1 - i)) = -\pi/4 + 2n\pi$ ,  $n$  an integer.




**Problem 3.**

(a) Write  $\frac{i}{2}$  in polar form.

**Solution:** Looking at the picture, we see  $|\frac{i}{2}| = \frac{1}{2}$ ,  $\text{Arg}\left(\frac{i}{2}\right) = \frac{\pi}{2}$ . So,  $\frac{i}{2} = \frac{1}{2}e^{i\pi/2}$ .




(b) Consider the diagram



Multiply the diagram by  $i/2$ , i.e., sketch the resulting image.

**Solution:** If  $z = re^{i\theta}$  then  $\frac{i}{2} \cdot z = \frac{1}{2} e^{i\pi/2} \cdot re^{i\theta} = \frac{r}{2} e^{i(\theta+\pi/2)}$ . So multiplication by  $i/2$  scales  $z$  by  $1/2$  and rotates it by  $\pi/2$  counterclockwise. The sketch is below.



**Problem 4.** Show  $\overline{e^{i\theta}} = e^{-i\theta}$ .

**Solution:**  $\overline{e^{i\theta}} = \overline{\cos \theta + i \sin \theta} = \cos \theta - i \sin \theta = \cos(-\theta) + i \sin(-\theta) = e^{-i\theta}$  ■


**Problem 5.** Compute  $(1 + \sqrt{3}i)^{10}$ . (Use polar form.)

**Solution:** We write  $1 + \sqrt{3}i$  in polar form.

Magnitude:  $|1 + \sqrt{3}i| = 2$ .

Argument: We recognize the  $30^\circ, 60^\circ, 90^\circ$  triangle. So,  $\text{Arg}(1 + \sqrt{3}i) = \pi/3$ .

Thus,  $1 + \sqrt{3}i = 2e^{i\pi/3}$ .



Thus,  $(1 + \sqrt{3}i)^{10} = (2e^{i\pi/3})^{10} = 2^{10} e^{i10\pi/3} = 2^{10} e^{i4\pi/3} = 2^{10} \cdot \left( \frac{-1 - \sqrt{3}i}{2} \right)$ .

(The step leading to  $2^{10} e^{i4\pi/3}$  is because  $10\pi/3 = 4\pi/3 + 2\pi$ .)

**Problem 6.** Compute  $I = \int e^x \cos(5x) dx$ .

**Solution:** Complexify:  $I_c = \int e^x e^{5xi} dx = \int e^{x(1+5i)} dx.$

Relation:  $I = \text{Re}(I_c).$

Computing:  $I_c = \frac{e^{x(1+5i)}}{1+5i}.$

We need to find the real part of  $I_c.$  In 18.03 we use polar form, so first we need to find the polar form of  $1+5i.$

$$|1+5i| = \sqrt{26}; \quad \boxed{\phi = \text{Arg}(1+5i) = \tan^{-1}(5) \text{ in Q1}}.$$

So,  $1+5i = \sqrt{26}e^{i\phi}.$  This gives,  $I_c = \frac{e^{(1+5i)x}}{1+5i} = \frac{e^x e^{5xi}}{\sqrt{26}e^{i\phi}} = \frac{e^x}{\sqrt{26}} e^{i(5x-\phi)}.$

Expanding this:  $I_c = \frac{e^x}{\sqrt{26}} (\cos(5x-\phi) + i \sin(5x-\phi)).$

Thus,  $\boxed{I = \text{Re}(I_c) = \frac{e^x}{\sqrt{26}} \cos(5x-\phi)}.$

Note: As we get comfortable with this, we will skip writing down so many of the steps.

### Problem 7.

(a) *Find the fifth roots of 1. Draw a picture.*


**Solution:** Write 1 in polar form with all possible  $\theta:$

$$1 = e^{i2\pi n}, \quad n \text{ any integer.}$$


We want  $z$  so that  $z^5 = 1 = e^{i2\pi n}.$  So,

$$z = e^{i2\pi n/5} = \underbrace{\widehat{e^0}, e^{i2\pi/5}, e^{i4\pi/5}, e^{i6\pi/5}, e^{i8\pi/5}}_{5 \text{ fifth roots}}, \overbrace{e^{i10\pi/5}}^{\text{same as } e^0 = 1}, \dots$$

On the plot, the roots are evenly spaced every  $2\pi/5 = 72^\circ$  around the unit circle.



Left: picture for Part (a)



right: picture for Part (b)

(b) *Find the fifth roots of  $1+i.$  Draw a picture.*

**Solution:** The idea is the same idea as Part (a). We have  $1+i = \sqrt{2}e^{i\pi/4+2n\pi}.$  So,  $z^5 = 1+i \Rightarrow z = 2^{1/10}e^{i(\pi/20+2n\pi/5)}.$  The 5 roots are


$$2^{1/10}e^{i\pi/20}, \quad 2^{1/10}e^{i9\pi/20}, \quad 2^{1/10}e^{i17\pi/20}, \quad 2^{1/10}e^{i25\pi/20}, \quad 2^{1/10}e^{i33\pi/20}.$$

Graphically the five roots are evenly spaced around the circle of radius  $2^{1/10}$ .


### Problem 8.

(a) Draw the trajectory of  $z = e^{it}$

**Solution:**  $z = e^{it} = \cos t + i \sin t$ . This always has unit length. So the trajectory goes round and round the unit circle.



Left: plot for Part (a);




right: plot for Part (b)

(b) Draw the trajectory of  $z = te^{it}$

**Solution:** This is similar to Part (a), except the factor of  $t$  causes the magnitude of  $z$  to grow. The trajectory spirals out from the origin.

(c) Plot the points  $e^{ij\pi/4}$ , for  $j = 0, 1, 2, 3 \dots$

**Solution:** Note:  $e^{i8\pi/4} = e^{i2\pi} = 1$ . These are the 8th roots of 1, i.e.,  $z^8 = 1$ .



### Problem 9.

(a) Write  $\sin t$  and  $\cos t$  in terms of  $e^{it}$  and  $e^{-it}$ .

**Solution:** Expand the exponentials: 
$$\begin{cases} e^{it} &= \cos t + i \sin t \\ e^{-it} &= \cos t - i \sin t \end{cases}$$

Adding and subtracting: 
$$\begin{cases} e^{it} + e^{-it} &= 2 \cos t \\ e^{it} - e^{-it} &= 2i \sin t \end{cases}$$
 So, 
$$\begin{cases} \cos t &= \frac{e^{it} + e^{-it}}{2} \\ \sin t &= \frac{e^{it} - e^{-it}}{2i} \end{cases}$$

(b) Find all real-valued functions of the form  $f(t) = c_1 e^{it} + c_2 e^{-it}$ , where  $c_1, c_2$  are complex constants.

**Solution:** From Part (a), we see that

$$\frac{1}{2}e^{it} + \frac{1}{2}e^{-it} = \cos t \quad \text{and} \quad \frac{1}{2i}e^{it} - \frac{1}{2i}e^{-it} = \sin t$$

are real-valued linear combinations of  $e^{it}$  and  $e^{-it}$ . Thus, for  $a, b$  real numbers,

$$\left(\frac{a}{2} + \frac{b}{2i}\right) e^{it} + \left(\frac{a}{2} - \frac{b}{2i}\right) e^{-it} = a \cos t + b \sin t$$

are the real-valued linear combinations of  $e^{it}$  and  $e^{-it}$ .

**Problem 10.** *Find all the roots of  $x^4 + x^2 = 0$ .*

**Solution:** This is fourth-order, so there are 4 roots. Factoring:

$$x^2(x^2 + 1) = 0 \quad \Rightarrow \quad x = 0, 0, i, -i \text{ are the 4 roots.}$$

MIT OpenCourseWare

<https://ocw.mit.edu>

ES.1803 Differential Equations

Spring 2024

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.